

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 November 2002 (28.11.2002)

PCT

(10) International Publication Number  
**WO 02/094823 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 471/04**,  
A61K 31/435, 31/495, A61P 11/06, 29/00

(21) International Application Number: PCT/CA02/00746

(22) International Filing Date: 22 May 2002 (22.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/293,247 24 May 2001 (24.05.2001) US

(71) Applicant (for all designated States except US):  
**MERCK FROSST CANADA & CO.** [CA/CA]; 16711  
Trans-Canada Highway, Kirkland, Québec H9H 3L1 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GUAY, Daniel**  
[CA/CA]; 16711 Trans-Canada Highway, Kirkland,  
Québec H9H 3L1 (CA). **GIRARD, Mario** [CA/CA];  
16711 Trans-Canada Highway, Kirkland, Québec H9H 3L1  
(CA). **HAMEL, Pierre** [CA/CA]; 16711 Trans-Canada  
Highway, Kirkland, Québec H9H 3L1 (CA). **LAL-  
IBERTE, Sebastien** [CA/CA]; 16711 Trans-Canada  
Highway, Kirkland, Québec H9H 3L1 (CA). **FRIESEN,  
Richard** [CA/CA]; 16711 Trans-Canada Highway, Kirk-  
land, Québec H9H 3L1 (CA). **GIRARD, Yves** [CA/CA];  
16711 Trans-Canada Highway, Kirkland, Québec H9H  
3L1 (CA). **LI, Chun** [CA/CA]; 16711 Trans-Canada  
Highway, Kirkland, Québec H9H 3L1 (CA).

(74) Agents: **OGILVY RENAULT** et al.; Attn: Murphy,  
Kevin, P., Suite 1600, 1981 McGill College Avenue,  
Montreal, Québec H3A 2Y3 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).


**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: 1-BIARYL-1,8-NAPHTHYRIDIN-4-ONE PHOSPHODIESTERASE-4 INHIBITORS

(57) Abstract: Compounds represented by Formula (I) or  
a pharmaceutically acceptable salt thereof, are phosphodi-  
esterase-4 inhibitors useful in the treatment of asthma and  
inflammation.



WO 02/094823 A1

mco23  
#2

## TITLE OF THE INVENTION

1-BIARYL-1,8-NAPHTHYRIDIN-4-ONE  
PHOSPHODIESTERASE-4 INHIBITORS

5

## BACKGROUND OF THE INVENTION

## FIELD OF THE INVENTION

The present invention is directed to compounds that are biaryl substituted 1,8-naphthyridin-4(1*H*)-ones. In particular, this invention is directed to phenyl or pyridyl substituted 1,8-naphthyridin-4(1*H*)-ones which are phosphodiesterase-4 inhibitors wherein the phenyl or pyridyl group is at the 1-position and contains an aryl substituent group further optionally substituted.

## 15 RELATED BACKGROUND

Hormones are compounds that variously affect cellular activity. In many respects, hormones act as messengers to trigger specific cellular responses and activities. Many effects produced by hormones, however, are not caused by the singular effect of just the hormone. Instead, the hormone first binds to a receptor, thereby triggering the release of a second compound that goes on to affect the cellular activity. In this scenario, the hormone is known as the first messenger while the second compound is called the second messenger. Cyclic adenosine monophosphate (adenosine 3', 5'-cyclic monophosphate, "cAMP" or "cyclic AMP") is known as a second messenger for hormones including epinephrine, glucagon, calcitonin, corticotrophin, lipotropin, luteinizing hormone, norepinephrine, parathyroid hormone, thyroid-stimulating hormone, and vasopressin. Thus, cAMP mediates cellular responses to hormones. Cyclic AMP also mediates cellular responses to various neurotransmitters.

Phosphodiesterases ("PDE") are a family of enzymes that metabolize 3', 5' cyclic nucleotides to 5' nucleoside monophosphates, thereby terminating cAMP second messenger activity. A particular phosphodiesterase, phosphodiesterase-4 ("PDE4", also known as "PDE-IV"), which is a high affinity, cAMP specific, type IV PDE, has generated interest as potential targets for the development of novel anti-asthmatic and anti-inflammatory compounds. PDE4 is known to exist as at least four

isoenzymes, each of which is encoded by a distinct gene. Each of the four known PDE4 gene products is believed to play varying roles in allergic and/or inflammatory responses. Thus, it is believed that inhibition of PDE4, particularly the specific PDE4 isoforms that produce detrimental responses, can beneficially affect allergy and inflammation symptoms. It would be desirable to provide novel compounds and compositions that inhibit PDE4 activity.

A major concern with the use of PDE4 inhibitors is the side effect of emesis which has been observed for several candidate compounds as described in C.Burnouf et al., ("Burnouf"), *Ann. Rep. In Med. Chem.*, 33:91-109(1998). B.Hughes et al., *Br. J.Pharmacol.*, 118:1183-1191(1996); M.J.Perry et al., *Cell Biochem. Biophys.*, 29:113-132(1998); S.B.Christensen et al., *J.Med. Chem.*, 41:821-835(1998); and Burnouf describe the wide variation of the severity of the undesirable side effects exhibited by various compounds. As described in M.D.Houslay et al., *Adv. In Pharmacol.*, 44:225-342(1998) and D.Spina et al., *Adv. In Pharmacol.*, 44:33-89(1998), there is great interest and research of therapeutic PDE4 inhibitors.

International Patent Publication WO9422852 describes quinolines as PDE4 inhibitors. International Patent Publication WO9907704 describes 1-aryl-1,8-naphthylidin-4-one derivatives as PDE4 inhibitors.

A.H.Cook, et al., *J.Chem. Soc.*, 413-417(1943) describes gamma-pyridylquinolines. Other quinoline compounds are described in Kei Manabe et al., *J.Org. Chem.*, 58(24):6692-6700(1993); Kei Manabe et al., *J.Am. Chem. Soc.*, 115(12):5324-5325(1993); and Kei Manabe et al., *J.Am. Chem. Soc.*, 114(17):6940-6941(1992).

Compounds that include ringed systems are described by various investigators as effective for a variety of therapies and utilities. For example, International Patent Publication No. WO 98/25883 describes ketobenzamides as calpain inhibitors, European Patent Publication No. EP 811610 and U.S. Patent Nos. 5,679,712, 5,693,672 and 5,747,541 describe substituted benzoylguanidine sodium channel blockers, U.S. Patent No. 5,736,297 describes ring systems useful as a photosensitive composition.


U.S. Patent Nos. 5,491,147, 5,608,070, 5,622,977, 5,739,144, 5,776,958, 5,780,477, 5,786,354, 5,798,373, 5,849,770, 5,859,034, 5,866,593, 5,891,896, and International Patent Publication WO 95/35283 describe PDE4 inhibitors that are tri-substituted aryl or heteroaryl phenyl derivatives. U.S. Patent No.

5,580,888 describes PDE4 inhibitors that are styryl derivatives. U.S. Patent No. 5,550,137 describes PDE4 inhibitors that are phenylaminocarbonyl derivatives. U.S. Patent No. 5,340,827 describes PDE4 inhibitors that are phenylcarboxamide compounds. U.S. Patent No. 5,780,478 describes PDE4 inhibitors that are tetra-substituted phenyl derivatives. International Patent Publication WO 96/00215 describes substituted oxime derivatives useful as PDE4 inhibitors. U.S. Patent No. 5,633,257 describes PDE4 inhibitors that are cyclo(alkyl and alkenyl)phenyl-alkenyl (aryl and heteroaryl) compounds.

However, there remains a need for novel compounds and compositions that therapeutically inhibit PDE4 with minimal side effects.

#### SUMMARY OF THE INVENTION

The present invention is directed to biaryl substituted 1,8-naphthyridin-4(1*H*)-ones represented by Formula (I):



(I)


or pharmaceutically acceptable salts thereof, which are phosphodiesterase-4 inhibitors.

This invention also provides a pharmaceutical composition which includes an effective amount of the novel biaryl substituted 1,8-naphthyridin-4(1*H*)-ones and a pharmaceutically acceptable carrier. This invention further provides a method of treatment in mammals of, for example, asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, ulcerative colitis,

Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, monopolar depression, acute and chronic neurodegenerative disorders with inflammatory components, Parkinson disease, Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour growth and cancerous invasion of normal tissues by the administration of an effective amount of the novel substituted 1,8-naphthyridin-4(1H)-ones or a precursor compound which forms *in vivo* the novel biaryl substituted 1,8-naphthyridin-4(1H)-ones which are phosphodiesterase-4 inhibitors.

#### DETAILED DESCRIPTION OF THE INVENTION

A compound of this invention is represented by Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein

Ar is phenyl, pyridyl, pyrimidyl, indolyl, quinolinyl, thienyl, pyridonyl, oxazolyl, oxadiazolyl, thiadiazolyl, or imidazolyl, or oxides thereof when Ar is a heteroaryl;

R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is H, or a -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkoxy, -C<sub>2-6</sub>alkenyl, -C<sub>3-6</sub>alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

In one aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is phenyl;

R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is H, or a -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkoxy, -C<sub>2-6</sub>alkenyl, -C<sub>3-6</sub>alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl,

-C(0-6alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

5 n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent  
10 halogen or -OH substituents.

In an embodiment of this one aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is phenyl;

15 R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is -C<sub>1-6</sub>alkyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any  
20 alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

25 n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent  
30 halogen or -OH substituents.

In another embodiment of this one aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

- Ar is phenyl;  
5 R is H or  $-C_{1-6}$ alkyl;  
R<sup>1</sup> is  $-C_{3-6}$ cycloalkyl, optionally substituted with 1-3 independent  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, OH, amino,  $-(C_{0-6}$ alkyl)-SO<sub>n</sub>-( $C_{1-6}$ alkyl), nitro, CN, =N-O- $C_{1-6}$ alkyl, -O-N= $C_{1-6}$ alkyl, or halogen substituents;  
R<sup>2</sup> is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkyl( $C_{3-6}$ cycloalkyl)( $C_{3-6}$ cycloalkyl),  $-C_{1-6}$ alkoxy, phenyl, heteroaryl, heterocycle, amino,  
10  $-C(O)-C_{1-6}$ alkyl,  $-C(O)-O-C_{1-6}$ alkyl,  $-C_{1-6}$ alkyl(=N-OH),  $-C(N=NOH)C_{1-6}$ alkyl,  $-C_{0-6}$ alkyl(oxy) $C_{1-6}$ alkyl-phenyl,  $-SO_nNH(C_{0-6}$ alkyl), or  $-(C_{0-6}$ alkyl)-SO<sub>n</sub>-( $C_{1-6}$ alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, hydroxy, amino, or  $-C(O)-O-C_{1-6}$ alkyl, and any  
15 alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;  
n is 0, 1, or 2;  
R<sup>3</sup> is H, OH, amine, halogen or  $C_{1-6}$ alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  
R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen,  $-C_{1-6}$ alkyl,  
20  $-C_{1-6}$ alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

In still another embodiment of this one aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof,  
25 wherein

- Ar is phenyl;  
R is H or  $-C_{1-6}$ alkyl;  
R<sup>1</sup> is pyridyl, optionally substituted with 1-3 independent  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, OH, amino,  $-(C_{0-6}$ alkyl)-SO<sub>n</sub>-( $C_{1-6}$ alkyl), nitro, CN, =N-O- $C_{1-6}$ alkyl, -O-N= $C_{1-6}$ alkyl, or halogen substituents;  
30 R<sup>2</sup> is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkyl( $C_{3-6}$ cycloalkyl)( $C_{3-6}$ cycloalkyl),  $-C_{1-6}$ alkoxy, phenyl, heteroaryl, heterocycle, amino,  $-C(O)-C_{1-6}$ alkyl,  $-C(O)-O-C_{1-6}$ alkyl,  $-C_{1-6}$ alkyl(=N-OH),  $-C(N=NOH)C_{1-6}$ alkyl,



-C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

5 n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent  
10 halogen or -OH substituents.

In a second aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridyl, pyrimidyl, or oxides thereof;

15 R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is H, or a -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkoxy, -C<sub>2-6</sub>alkenyl, -C<sub>3-6</sub>alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -  
(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or  
20 halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with  
25 halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  
30 R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

In an embodiment of the second aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridyl, pyrimidyl, or oxides thereof;

5 R is H;

R<sup>1</sup> is H;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, 10 -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

15 R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

20

In another embodiment of the second aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridyl, pyrimidyl, or oxides thereof;

25 R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is -C<sub>1-6</sub>alkyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, 30 -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with

halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is

5 optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

10 In still another embodiment of the second aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridyl, pyrimidyl, or oxides thereof;

R is H or -C<sub>1-6</sub>alkyl;

15 R<sup>1</sup> is -C<sub>3-6</sub>cycloalkyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

20 R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

25 n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

30 R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

In yet another embodiment of the second aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

- Ar is pyridyl, pyrimidyl, or oxides thereof;  
R is H or -C<sub>1-6</sub>alkyl;  
R<sup>1</sup> is pyridyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;  
R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;  
n is 0, 1, or 2;  
R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  
R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

- In a third aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

- Ar is indolyl, quinolinyl, or oxides thereof;  
R is H or -C<sub>1-6</sub>alkyl;  
R<sup>1</sup> is H, or a -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkoxy, -C<sub>2-6</sub>alkenyl, -C<sub>3-6</sub>alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

- R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;
- n is 0, 1, or 2;
- R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and
- R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

- 15 In an embodiment of the third aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein
- Ar is indolyl;
- R is H or -C<sub>1-6</sub>alkyl;
- R<sup>1</sup> is -C<sub>1-6</sub>alkyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;
- 20 R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;
- 25 n is 0, 1, or 2;
- 30 R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

$R^4$ ,  $R^5$ ,  $R^6$ , and  $R^7$  each independently is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents.

5 In another embodiment of the third aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is quinolinyl or oxide thereof;

R is H or  $-C_{1-6}$ alkyl;

10  $R^1$  is  $-C_{1-6}$ alkyl, optionally substituted with 1-3 independent  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, OH, amino,  $-(C_{0-6}$ alkyl) $-SO_n-(C_{1-6}$ alkyl), nitro, CN,  $=N-O-C_{1-6}$ alkyl,  $-O-N=C_{1-6}$ alkyl, or halogen substituents;

$R^2$  is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkyl( $C_{3-6}$ cycloalkyl)( $C_{3-6}$ cycloalkyl),  $-C_{1-6}$ alkoxy, phenyl, heteroaryl, heterocycle, amino, 15  $-C(O)-C_{1-6}$ alkyl,  $-C(O)-O-C_{1-6}$ alkyl,  $-C_{1-6}$ alkyl( $=N-OH$ ),  $-C(N=NOH)C_{1-6}$ alkyl,  $-C_{0-6}$ alkyl(oxy) $C_{1-6}$ alkyl-phenyl,  $-SO_nNH(C_{0-6}$ alkyl), or  $-(C_{0-6}$ alkyl) $-SO_n-(C_{1-6}$ alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, hydroxy, amino, or  $-C(O)-O-C_{1-6}$ alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents;

20 n is 0, 1, or 2;

$R^3$  is H, OH, amine, halogen or  $C_{1-6}$ alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

$R^4$ ,  $R^5$ ,  $R^6$ , and  $R^7$  each independently is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent 25 halogen or  $-OH$  substituents.

In a fourth aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

30 Ar is thienyl;

R is H or  $-C_{1-6}$ alkyl;

$R^1$  is H, or a  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkoxy,  $-C_{2-6}$ alkenyl,  $-C_{3-6}$ alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is

optionally substituted with 1-3 independent  $-C_1$ -6alkyl,  $-C_1$ -6alkoxy, OH, amino,  $-(C_0$ -6alkyl)- $SO_n$ -( $C_1$ -6alkyl), nitro, CN,  $=N-O-C_1$ -6alkyl,  $-O-N=C_1$ -6alkyl, or halogen substituents;

- 5  $R^2$  is H, halogen,  $-C_1$ -6alkyl,  $-C_3$ -6cycloalkyl,  $-C_1$ -6alkyl( $C_3$ -6cycloalkyl)( $C_3$ -6cycloalkyl),  $-C_1$ -6alkoxy, phenyl, heteroaryl, heterocycle, amino,  $-C(O)-C_1$ -6alkyl,  $-C(O)-O-C_1$ -6alkyl,  $-C_1$ -6alkyl( $=N-OH$ ),  $-C(N=NOH)C_1$ -6alkyl,  $-C_0$ -6alkyl(oxy) $C_1$ -6alkyl-phenyl,  $-SO_nNH(C_0$ -6alkyl), or  $-(C_0$ -6alkyl)- $SO_n$ -( $C_1$ -6alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen,  $-C_1$ -6alkyl,  $-C_1$ -6alkoxy, hydroxy, amino, or  $-C(O)-O-C_1$ -6alkyl, and any
- 10 alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents;
- $n$  is 0, 1, or 2;
- $R^3$  is H, OH, amine, halogen amine, halogen or  $C_1$ -6alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and
- 15  $R^4$ ,  $R^5$ ,  $R^6$ , and  $R^7$  each independently is H, halogen,  $-C_1$ -6alkyl,  $-C_1$ -6alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents.

- 20 In an embodiment of the fourth aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

- Ar is thienyl;
- R is H or  $-C_1$ -6alkyl;
- $R^1$  is  $-C_3$ -6cycloalkyl, optionally substituted with 1-3 independent  $-C_1$ -6alkyl,  $-C_1$ -6alkoxy, OH, amino,  $-(C_0$ -6alkyl)- $SO_n$ -( $C_1$ -6alkyl), nitro, CN,  $=N-O-C_1$ -6alkyl,  $-O-N=C_1$ -6alkyl, or halogen substituents;
- 25  $R^2$  is H, halogen,  $-C_1$ -6alkyl,  $-C_3$ -6cycloalkyl,  $-C_1$ -6alkyl( $C_3$ -6cycloalkyl)( $C_3$ -6cycloalkyl),  $-C_1$ -6alkoxy, phenyl, heteroaryl, heterocycle, amino,  $-C(O)-C_1$ -6alkyl,  $-C(O)-O-C_1$ -6alkyl,  $-C_1$ -6alkyl( $=N-OH$ ),  $-C(N=NOH)C_1$ -6alkyl,  $-C_0$ -6alkyl(oxy) $C_1$ -6alkyl-phenyl,  $-SO_nNH(C_0$ -6alkyl), or  $-(C_0$ -6alkyl)- $SO_n$ -( $C_1$ -6alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen,  $-C_1$ -6alkyl,  $-C_1$ -6alkoxy, hydroxy, amino, or  $-C(O)-O-C_1$ -6alkyl, and any
- 30 alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents;
- $n$  is 0, 1, or 2;

$R^3$  is H, OH, amine, halogen or  $C_{1-6}$ alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  $R^4$ ,  $R^5$ ,  $R^6$ , and  $R^7$  each independently is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents.

In a fifth aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridonyl;

R is H or  $-C_{1-6}$ alkyl;

$R^1$  is H, or a  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkoxy,  $-C_{2-6}$ alkenyl,  $-C_{3-6}$ alkynyl, heteroaryl, or heterocycle group, wherein any of the groups is optionally substituted with 1-3 independent  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, OH, amino,  $-(C_{0-6}$ alkyl)- $SO_n$ -( $C_{1-6}$ alkyl), nitro, CN,  $=N-O-C_{1-6}$ alkyl,  $-O-N=C_{1-6}$ alkyl, or halogen substituents;

$R^2$  is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{3-6}$ cycloalkyl,  $-C_{1-6}$ alkyl( $C_{3-6}$ cycloalkyl)( $C_{3-6}$ cycloalkyl),  $-C_{1-6}$ alkoxy, phenyl, heteroaryl, heterocycle, amino,  $-C(O)-C_{1-6}$ alkyl,  $-C(O)-O-C_{1-6}$ alkyl,  $-C_{1-6}$ alkyl( $=N-OH$ ),  $-C(N=NOH)C_{1-6}$ alkyl,  $-C_{0-6}$ alkyl(oxy) $C_{1-6}$ alkyl-phenyl,  $-SO_nNH(C_{0-6}$ alkyl), or  $-(C_{0-6}$ alkyl)- $SO_n$ -( $C_{1-6}$ alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, hydroxy, amino, or  $-C(O)-O-C_{1-6}$ alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents;

n is 0, 1, or 2;

$R^3$  is H, OH, amine, halogen or  $C_{1-6}$ alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  $R^4$ ,  $R^5$ ,  $R^6$ , and  $R^7$  each independently is H, halogen,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or  $-OH$  substituents.

In an embodiment of the fifth aspect, a compound of this invention is represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

Ar is pyridonyl;



R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is -C<sub>3-6</sub>cycloalkyl, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

5 R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with  
10 halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and  
15 R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

20 As used herein, "alkyl" as well as other groups having the prefix "alk" such as, for example, alkoxy, alkanoyl, alkenyl, alkynyl and the like, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl and the like. "Alkenyl", "alkynyl" and other like terms include carbon  
25 chains containing at least one unsaturated C-C bond.

The term "cycloalkyl" means carbocycles containing no heteroatoms, and includes mono-, bi- and tricyclic saturated carbocycles, as well as fused ring systems. Such fused ring systems can include one ring that is partially or fully unsaturated such as a benzene ring to form fused ring systems such as benzofused  
30 carbocycles. Cycloalkyl includes such fused ring systems as spirofused ring systems. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, decahydronaphthalene, adamantane, indanyl, indenyl, fluorenyl, 1,2,3,4-tetrahydronaphthalene and the like. Similarly, "cycloalkenyl" means carbocycles

containing no heteroatoms and at least one non-aromatic C-C double bond, and include mono-, bi- and tricyclic partially saturated carbocycles, as well as benzofused cycloalkenes. Examples of cycloalkenyl include cyclohexenyl, indenyl, and the like.

5 The term "cycloalkyloxy" unless specifically stated otherwise includes a cycloalkyl group connected to the oxy connecting atom.

The term "alkoxy" unless specifically stated otherwise includes an alkyl group connected to the oxy connecting atom.

10 The term "aryl" unless specifically stated otherwise includes multiple ring systems as well as single ring systems such as, for example, phenyl or naphthyl. The term "aryloxy" unless specifically stated otherwise includes multiple ring systems as well as single ring systems such as, for example, phenyl or naphthyl, connected through the oxy connecting atom to the connecting site.

15 The term "C<sub>0</sub>-C<sub>6</sub>alkyl" includes alkyls containing 6, 5, 4, 3, 2, 1, or no carbon atoms. An alkyl with no carbon atoms is a hydrogen atom substituent or a direct bond – depending on whether the alkyl is a terminus or a bridging moiety.

20 The term "hetero" unless specifically stated otherwise includes one or more O, S, or N atoms. For example, heterocycloalkyl and heteroaryl include ring systems that contain one or more O, S, or N atoms in the ring, including mixtures of such atoms. The hetero atoms replace ring carbon atoms. Thus, for example, a heterocycloC<sub>5</sub>alkyl is a five membered ring containing from 5 to no carbon atoms.

25 Examples of heteroaryl include, for example, pyridinyl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinoxalinyl, furyl, benzofuryl, dibenzofuryl, thienyl, benzothienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl.

The term "heteroaryloxy" unless specifically stated otherwise describes a heteroaryl group connected through an oxy connecting atom to the connecting site.

30 Examples of heteroaryl(C<sub>1-6</sub>)alkyl include, for example, furylmethyl, furylethyl, thienylmethyl, thienylethyl, pyrazolylmethyl, oxazolylmethyl, oxazolylethyl, isoxazolylmethyl, thiazolylmethyl, thiazolylethyl, imidazolylmethyl, imidazolylethyl, benzimidazolylmethyl, oxadiazolylmethyl, oxadiazolylethyl, thiadiazolylmethyl, thiadiazolylethyl, triazolylmethyl, triazolylethyl, tetrazolylmethyl, tetrazolylethyl, pyridinylmethyl, pyridinylethyl, pyridazinylmethyl,

pyrimidinylmethyl, pyrazinylmethyl, quinolinylmethyl, isoquinolinylmethyl and quinoxalinylmethyl.

Examples of heterocycloC<sub>3-7</sub>alkyl include, for example, azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, imidazoliny, pyrolidin-2-one, piperidin-2-one, and thiomorpholinyl.

Examples of aryl(C<sub>1-6</sub>)alkyl include, for example, phenyl(C<sub>1-6</sub>)alkyl, and naphthyl(C<sub>1-6</sub>)alkyl.

Examples of heterocycloC<sub>3-7</sub>alkylcarbonyl(C<sub>1-6</sub>)alkyl include, for example, azetidiny carbonyl(C<sub>1-6</sub>)alkyl, pyrrolidinyl carbonyl(C<sub>1-6</sub>)alkyl, piperidinyl carbonyl(C<sub>1-6</sub>)alkyl, piperazinyl carbonyl(C<sub>1-6</sub>)alkyl, morpholinyl carbonyl(C<sub>1-6</sub>)alkyl, and thiomorpholinyl carbonyl(C<sub>1-6</sub>)alkyl.

The term "amine" unless specifically stated otherwise includes primary, secondary and tertiary amines.

Unless otherwise stated, the term "carbamoyl" is used to include -NHC(O)OC<sub>1-4</sub>alkyl, and -OC(O)NHC<sub>1-4</sub>alkyl.

The term "halogen" includes fluorine, chlorine, bromine and iodine atoms.

The term "optionally substituted" is intended to include both substituted and unsubstituted. Thus, for example, optionally substituted aryl could represent a pentafluorophenyl or a phenyl ring. Further, the substitution can be made at any of the groups. For example, substituted aryl(C<sub>1-6</sub>)alkyl includes substitution on the aryl group as well as substitution on the alkyl group.

The term "oxide" of heteroaryl groups is used in the ordinary well-known chemical sense and include, for example, *N*-oxides of nitrogen heteroatoms.

Compounds described herein contain one or more double bonds and may thus give rise to cis/trans isomers as well as other conformational isomers. The present invention includes all such possible isomers as well as mixtures of such isomers.

Compounds described herein can contain one or more asymmetric centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and pharmaceutically acceptable salts thereof. The above Formula I is shown without a definitive stereochemistry at certain positions. The present invention includes all

stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Further, mixtures of stereoisomers as well as isolated specific stereoisomers are also included. During the course of the synthetic procedures used to prepare such compounds, or in using racemization or epimerization procedures known to those skilled in the art, the products of such procedures can be a mixture of stereoisomers.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (ic and ous), ferric, ferrous, lithium, magnesium, manganese (ic and ous), potassium, sodium, zinc and the like salts. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, as well as cyclic amines and substituted amines such as naturally occurring and synthesized substituted amines. Other pharmaceutically acceptable organic non-toxic bases from which salts can be formed include ion exchange resins such as, for example, arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

When the compound of the present invention is basic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Particularly preferred are benzenesulfonic, citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (or pharmaceutically acceptable salts thereof) as an active ingredient, a pharmaceutically acceptable carrier and optionally other therapeutic ingredients or adjuvants. Such additional therapeutic ingredients include, for example, i) Leukotriene receptor antagonists, ii) Leukotriene biosynthesis inhibitors, iii) corticosteroids, iv) H1 receptor antagonists, v) beta 2 adrenoceptor agonists, vi) COX-2 selective inhibitors, vii) statins, viii) non-steroidal anti-inflammatory drugs ("NSAID"), and ix) M2/M3 antagonists. The compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

Creams, ointments, jellies, solutions, or suspensions containing the compound of Formula I can be employed for topical use. Mouth washes and gargles are included within the scope of topical use for the purposes of this invention.

Dosage levels from about 0.001mg/kg to about 140mg/kg of body weight per day are useful in the treatment of conditions such as asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, monopolar depression, acute and chronic neurodegenerative disorders with inflammatory components, Parkinson disease, Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour

growth and cancerous invasion of normal tissues which are responsive to PDE4 inhibition, or alternatively about 0.05mg to about 7g per patient per day. For example, inflammation may be effectively treated by the administration of from about 0.01mg to 50mg of the compound per kilogram of body weight per day, or  
5 alternatively about 0.5mg to about 2.5g per patient per day. Further, it is understood that the PDE4 inhibiting compounds of this invention can be administered at prophylactically effective dosage levels to prevent the above-recited conditions.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated  
10 and the particular mode of administration. For example, a formulation intended for the oral administration to humans may conveniently contain from about 0.5mg to about 5g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms will generally contain between from about 0.01mg to  
15 about 1000mg of the active ingredient, typically 0.01mg, 0.05mg, 0.25mg, 1mg, 5mg, 25mg, 50mg, 100mg, 200mg, 300mg, 400mg, 500mg, 600mg, 800mg or 1000mg.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion,  
20 drug combination and the severity of the particular disease undergoing therapy.

In practice, the compounds represented by Formula I, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide  
25 variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient. Further, the compositions can be presented as a powder, as  
30 granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid, as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compound represented by Formula I, or pharmaceutically acceptable salts thereof, may also be administered by controlled release means and/or delivery devices. The compositions may be prepared by any of

the methods of pharmacy. In general, such methods include a step of bringing into association the active ingredient with the carrier that constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

Thus, the pharmaceutical compositions of this invention may include a pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable salt of Formula I. The compounds of Formula I, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

The pharmaceutical carrier employed can be, for example, a solid, liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

In preparing the compositions for oral dosage form, any convenient pharmaceutical media may be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like may be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like may be used to form oral solid preparations such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques

A tablet containing the composition of this invention may be prepared by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Each tablet preferably contains from about 0.1mg to about 500mg of the active ingredient and

each cachet or capsule preferably containing from about 0.1mg to about 500mg of the active ingredient.

Pharmaceutical compositions of the present invention suitable for parenteral administration may be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, 5 hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

Pharmaceutical compositions of the present invention suitable for 10 injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability. The pharmaceutical compositions must be stable under the conditions of manufacture 15 and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

Pharmaceutical compositions of the present invention can be in a form 20 suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, or the like. Further, the compositions can be in a form suitable for use in transdermal devices. These formulations may be prepared, utilizing a compound represented by Formula I of this invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or 25 ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the 30 mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.



In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above may include, as appropriate, one or more additional carrier ingredients such as diluents, buffers, flavoring agents, binders, surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation isotonic with the blood of the intended recipient. Compositions containing a compound described by Formula I, or pharmaceutically acceptable salts thereof, may also be prepared in powder or liquid concentrate form.

The compounds and pharmaceutical compositions of this invention have been found to exhibit biological activity as PDE4 inhibitors. Accordingly, another aspect of the invention is the treatment in mammals of, for example, asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, monopolar depression, acute and chronic neurodegenerative disorders with inflammatory components, Parkinson disease, Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour growth and cancerous invasion of normal tissues – maladies that are amenable to amelioration through inhibition of the PDE4 isoenzyme and the resulting elevated cAMP levels – by the administration of an effective amount of the compounds of this invention. The term "mammals" includes humans, as well as other animals such as, for example, dogs, cats, horses, pigs, and cattle. Accordingly, it is understood that the treatment of mammals other than humans is the treatment of clinical correlating afflictions to those above recited examples that are human afflictions.

- Further, as described above, the compound of this invention can be utilized in combination with other therapeutic compounds. In particular, the combinations of the PDE4 inhibiting compound of this invention can be advantageously used in combination with i) Leukotriene receptor antagonists, ii) Leukotriene biosynthesis inhibitors, iii) COX-2 selective inhibitors, iv) statins, v) NSAIDs, vi) M2/M3 antagonists, vii) corticosteroids, viii) H1 (histamine) receptor antagonists and ix) beta 2 adrenoceptor agonist.

- The abbreviations used herein have the following tabulated meanings. Abbreviations not tabulated below have their meanings as commonly used unless specifically stated otherwise.

Ac	=	acetyl
Bn	=	benzyl
CAMP		cyclic adenosine-3',5'-monophosphate
DBU	=	1,8-diazabicyclo[5.4.0]undec-7-ene
DIBAL	=	diisobutylaluminum hydride
DMAP	=	4-(dimethylamino)pyridine
DMF	=	N,N-dimethylformamide
Et <sub>3</sub> N	=	triethylamine
GST		glutathione transferase
HMDS		hexamethyldisilazide
LDA	=	lithium diisopropylamide
m-CPBA	=	metachloroperbenzoic acid
MMPP	=	monoperoxyphthalic acid
MPPM	=	monoperoxyphthalic acid, magnesium salt 6H <sub>2</sub> O
Ms	=	methanesulfonyl = mesyl = SO <sub>2</sub> Me
MsO	=	methanesulfonate = mesylate
NSAID	=	non-steroidal anti-inflammatory drug
o-Tol	=	ortho-tolyl
OXONE®	=	2KHSO <sub>5</sub> •KHSO <sub>4</sub> •K <sub>2</sub> SO <sub>4</sub>
PCC	=	pyridinium chlorochromate
PDC	=	pyridinium dichromate

PDE		phosphodiesterase
Ph	=	phenyl
Phe	=	benzenediyl
PMB	=	para-methoxybenzyl
Pye	=	pyridinediyl
r.t.	=	room temperature
Rac.	=	racemic
SAM	=	aminosulfonyl or sulfonamide or SO <sub>2</sub> NH <sub>2</sub>
SEM	=	2-(trimethylsilyl)ethoxymethoxy
SPA	=	scintillation proximity assay
TBAF	=	tetra-n-butylammonium fluoride
Th	=	2- or 3-thienyl
TFA	=	trifluoroacetic acid
TFAA	=	trifluoroacetic acid anhydride
THF	=	tetrahydrofuran
Thi	=	thiophenediyl
TLC	=	thin layer chromatography
TMS-CN	=	trimethylsilyl cyanide
TMSI		trimethylsilyl iodide
Tz	=	1H (or 2H)-tetrazol-5-yl
CAN		ceric ammonium nitrate
C <sub>3</sub> H <sub>5</sub>	=	allyl

#### ALKYL GROUP ABBREVIATIONS

Me	=	Methyl
Et	=	ethyl
<i>n</i> -Pr	=	normal propyl
<i>i</i> -Pr	=	isopropyl
<i>n</i> -Bu	=	normal butyl
<i>i</i> -Bu	=	isobutyl
<i>s</i> -Bu	=	secondary butyl

<i>t</i> -Bu	=	tertiary butyl
c-Pr	=	cyclopropyl
c-Bu	=	Cyclobutyl
c-Pen	=	cyclopentyl
c-Hex	=	cyclohexyl

## ASSAYS DEMONSTRATING BIOLOGICAL ACTIVITY

5

### LPS AND FMLP-INDUCED TNF- $\alpha$ AND LTB<sub>4</sub> ASSAYS IN HUMAN WHOLE BLOOD

Whole blood provides a protein and cell-rich milieu appropriate for the study of biochemical efficacy of anti-inflammatory compounds such as PDE4-selective inhibitors. Normal non-stimulated human blood does not contain detectable levels of TNF- $\alpha$  and LTB<sub>4</sub>. Upon stimulation with LPS, activated monocytes express and secrete TNF- $\alpha$  up to 8 hours and plasma levels remain stable for 24 hours. Published studies have shown that inhibition of TNF- $\alpha$  by increasing intracellular cAMP via PDE4 inhibition and/or enhanced adenylyl cyclase activity occurs at the transcriptional level. LTB<sub>4</sub> synthesis is also sensitive to levels of intracellular cAMP and can be completely inhibited by PDE4-selective inhibitors. As there is little LTB<sub>4</sub> produced during a 24 hour LPS stimulation of whole blood, an additional LPS stimulation followed by FMLP challenge of human whole blood is necessary for LTB<sub>4</sub> synthesis by activated neutrophils. Thus, by using the same blood sample, it is possible to evaluate the potency of a compound on two surrogate markers of PDE4 activity in the whole blood by the following procedure.

Fresh blood was collected in heparinized tubes by venipuncture from healthy human volunteers (male and female). These subjects had no apparent inflammatory conditions and had not taken any NSAIDs for at least 4 days prior to blood collection. 500 $\mu$ L aliquots of blood were pre-incubated with either 2 $\mu$ L of vehicle (DMSO) or 2 $\mu$ L of test compound at varying concentrations for 15 minutes at 37°C. This was followed by the addition of either 10 $\mu$ L vehicle (PBS) as blanks or 10 $\mu$ L LPS (1 $\mu$ g/mL final concentration, #L-2630 (Sigma Chemical Co., St. Louis,

MO) from *E. coli*, serotype 0111:B4; diluted in 0.1% w/v BSA (in PBS)). After 24 hours of incubation at 37°C; another 10 µL of PBS (blank) or 10 µL of LPS (1 µg/mL final concentration) was added to blood and incubated for 30 minutes at 37°C. The blood was then challenged with either 10 µL of PBS (blank) or 10 µL of fMLP (1 µM final concentration, #F-3506 (Sigma); diluted in 1% w/v BSA (in PBS)) for 15 minutes at 37°C. The blood samples were centrifuged at 1500xg for 10 minutes at 4°C to obtain plasma. A 50 µL aliquot of plasma was mixed with 200 µL methanol for protein precipitation and centrifuged as above. The supernatant was assayed for LTB<sub>4</sub> using an enzyme immunoassay kit (#520111 from Cayman Chemical Co., Ann Arbor, MI) according to the manufacturer's procedure. TNF-α was assayed in diluted plasma (in PBS) using an ELISA kit (Cistron Biotechnology, Pine Brook, NJ) according to manufacturer's procedure. The IC<sub>50</sub> values of Examples 1 to 76 generally ranged from 0.005 µM to 15.4 µM.

15

#### ANTI-ALLERGIC ACTIVITY *IN VIVO*

Compounds of the invention have been tested for effects on an IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs. Guinea pigs were initially sensitized to ovalbumin under mild cyclophosphamide-induced immunosuppression, by intraperitoneal injection of antigen in combinations with aluminum hydroxide and pertussis vaccine. Booster doses of antigen were given two and four weeks later. At six weeks, animals were challenged with aerosolized ovalbumin while under cover of an intraperitoneally administered anti-histamine agent (mepyramine). After a further 48h, bronchial alveolar lavages (BAL) were performed and the numbers of eosinophils and other leukocytes in the BAL fluids were counted. The lungs were also removed for histological examination for inflammatory damage. Administration of compounds of the Examples (0.001-10mg/kg i.p. or p.o.), up to three times during the 48h following antigen challenge, lead to a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes. There was also less inflammatory damage in the lungs of animals treated with compounds of the Examples.

30

#### SPA BASED PDE ACTIVITY ASSAY PROTOCOL

Compounds which inhibit the hydrolysis of cAMP to AMP by the type-IV cAMP-specific phosphodiesterases were screened in a 96-well plate format as follows:

In a 96 well-plate at 30°C was added the test compound (dissolved in 2μL DMSO), 188μL of substrate buffer containing [2,8-<sup>3</sup>H] adenosine 3',5'-cyclic phosphate (cAMP, 100nM to 50μM), 10mM MgCl<sub>2</sub>, 1mM EDTA, 50mM Tris, pH 7.5. The reaction was initiated by the addition of 10μL of human recombinant PDE4 (the amount was controlled so that ~10% product was formed in 10min.). The reaction was stopped after 10min. by the addition of 1mg of PDE-SPA beads (Amersham Pharmacia Biotech, Inc., Piscataway, NJ). The product AMP generated was quantified on a Wallac Microbeta® 96-well plate counter (EG&G Wallac Co., Gaithersburg, MD). The signal in the absence of enzyme was defined as the background. 100% activity was defined as the signal detected in the presence of enzyme and DMSO with the background subtracted. Percentage of inhibition was calculated accordingly. IC<sub>50</sub> value was approximated with a non-linear regression fit using the standard 4-parameter/multiple binding sites equation from a ten point titration.

The IC<sub>50</sub> values of Examples 1 to 76 were determined with 100nM cAMP using the purified GST fusion protein of the human recombinant phosphodiesterase IVa (met-248) produced from a baculovirus/Sf-9 expression system. The IC<sub>50</sub> values of Examples 1-76 generally ranged from 0.1 nM to 14.8 nM, although six examples had an IC<sub>50</sub> value between 34.3 and 134.0 nM.

The examples that follow are intended as an illustration of certain preferred embodiments of the invention and no limitation of the invention is implied.

Unless specifically stated otherwise, the experimental procedures were performed under the following conditions. All operations were carried out at room or ambient temperature - that is, at a temperature in the range of 18-25°C. Evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000pascals: 4.5-30mm Hg) with a bath temperature of up to 60°C. The course of reactions was followed by thin layer chromatography (TLC) and reaction times are

given for illustration only. Melting points are uncorrected and "d" indicates decomposition. The melting points given are those obtained for the materials prepared as described. Polymorphism may result in isolation of materials with different melting points in some preparations. The structure and purity of all final products were assured by at least one of the following techniques: TLC, mass spectrometry, nuclear magnetic resonance (NMR) spectrometry or microanalytical data. When given, yields are for illustration only. When given, NMR data is in the form of delta ( $\delta$ ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, determined at 300 MHz, 400 MHz or 500 MHz using the indicated solvent. Conventional abbreviations used for signal shape are: s. singlet; d. doublet; t. triplet; m. multiplet; br. broad; etc. In addition, "Ar" signifies an aromatic signal. Chemical symbols have their usual meanings; the following abbreviations have also been used: v (volume), w (weight), b.p. (boiling point), m.p. (melting point), L (liter(s)), mL (milliliters), g (gram(s)), mg (milligrams(s)), mol (moles), mmol (millimoles), eq (equivalent(s)).

### Methods of Synthesis

Compounds of the present invention can be prepared according to the following methods. The substituents are the same as in Formula I except where defined otherwise.

### Scheme 1

In a first method outlined in **Scheme 1** below, an appropriately substituted derivative of ethyl 2-chloronicotinoyl acetate of formula **II** is reacted with 1.5 equivalents of triethyl orthoformate and 5 equivalents of acetic anhydride at 130°C, and after removal of the volatile components, the crude 2-chloronicotinoyl acrylate of formula **III** is immediately reacted with 1.2 equivalents of an appropriately substituted haloaryl amine of formula **IV**, such as, for example 3-bromoaniline, in a halogenated hydrocarbon solvent such as methylene chloride at a temperature of 0°C to room temperature. After an appropriate reaction time ranging from 2 to 24 hours

the resulting 3-arylamino acrylate of formula **V** is obtained by evaporation of the solvent and may be further purified by chromatography on silica gel or crystallization from an appropriate solvent.

5                   The compound of formula **V** may alternatively be used without further purification in the following step. Cyclization of the compound of formula **V** to the 1-haloaryl-1,4-dihydro[1,8]naphthyridin-4-one carboxylate of formula **VI** is effected by treatment with a small excess of a strong base such as an alkali metal hydride, for example sodium hydride, in an appropriate solvent such as tetrahydrofuran at a  
10                   starting temperature of 0°C with warming to room temperature if required to complete the process. The product of formula **VI** is isolated in crude form by dilution with a large volume of water followed by filtration or by extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified  
15                   by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

                  The product of formula **VI** thus obtained can be hydrolyzed to the corresponding carboxylic acid derivative under basic conditions, using an aqueous  
20                   solution of an alkali base such as an alkali carbonate or preferably sodium or potassium hydroxide, with an organic cosolvent such as tetrahydrofuran or a primary, secondary or tertiary alkanol, such as methanol or ethanol, or a combination thereof at temperatures ranging from room temperature to reflux temperature for the appropriate time. The resultant carboxylic acid is isolated in crude form following acidification  
25                   using an aqueous solution of an inorganic acid such as hydrochloric, sulfuric or a similar acid, and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by  
30                   filtration.

                  The carboxylic acid is then transformed into the appropriate primary, secondary or tertiary amide analog of formula **VII** by any general procedure well



known to the organic chemist, preferably via initial transformation into a mixed anhydride by treatment with a small excess, such as 1.25 equivalents, of an appropriate alkyl chloroformate such as ethyl or isobutyl chloroformate, in the presence of a larger excess, such as 2.5 equivalents, of a tertiary organic amine such as triethylamine or N,N-diisopropylethylamine in an organic solvent such as tetrahydrofuran at low temperature, preferably 0°C, for a period of 30 minutes to 3 hours. An excess, usually 5 or more equivalents, of an appropriate primary or secondary amine or of an aqueous solution of ammonium hydroxide is then added and the resulting reaction is allowed to proceed at a temperature ranging from 0°C to room temperature for an appropriate length of time, usually 1-24 hours.

The desired amide of formula VII is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration. In cases where the amide moiety is 2,6-dichloropyridin-4-yl, a different procedure is used in which the anion of 4-amino-3,5-dichloropyridine is generated at low temperature, preferably at 0°C, using a strong alkali hydride such as sodium hydride in a solvent such as tetrahydrofuran, and reacted with the acid chloride of a carboxylic acid (from hydrolysis of an ester of formula VI) generated by an appropriate known procedure, usually by the action of oxalyl chloride activated by a catalytic amount of N,N-dimethylformamide in a solvent such as tetrahydrofuran.


The amides of general formula VII are processed into the products of formula I by reaction with an appropriately substituted aryl or heteroaryl boronic acid or boronate ester of formula VIII under the catalysis of a transition metal species such as *trans*-dibromobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in an appropriate solvent or solvent mixture, preferably a 1:1 mixture of toluene and ethanol in the presence of an excess of an aqueous solution of an alkali base such as sodium carbonate, at an appropriate temperature, preferably 50 to 100°C, for an appropriate length of time ranging from 0.5 to 48 hours.

The resulting reaction product is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

Compounds of formula I may also be obtained by reaction of a compound of formula VII with an appropriately substituted aryl or heteroaryl tributyl stannane of formula IX under the catalysis of a transition metal species such as *trans*-dibromobis(triphenylphosphine)palladium (II) or [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in the presence of a copper (I) species such as cuprous iodide in an appropriate solvent such as N,N-dimethylformamide at a temperature range of 50-100°C for a period of 2 to 24 hours. Isolation of the reaction product is effected as described above.

Alternatively, an ester of formula VI can be processed into an ester of formula X by reaction with an appropriately substituted boronic acid or boronate ester, or with an appropriately substituted stannane derivative under the conditions described above, and the ester can be hydrolyzed and transformed into an amide of formula I.

The boronic acids of formula VIII or corresponding boronate esters are usually obtained from commercial sources. Where required, they can be prepared readily from the corresponding halides via metallation with n-butyllithium followed by reaction with a trialkyl borate, or by using classical transition metal-catalyzed coupling procedures using diboron pinacol ester. The stannanes of formula IX are generated from the corresponding halides via initial metallation using n-butyllithium followed by addition of tributyltin chloride.




## Scheme 2

In an alternative method for the preparation of compounds of formula I, outlined in **Scheme 2** below, an amide of formula VII can be transformed into a  
5 corresponding boronate ester of formula XI by treatment with an excess of diboron pinacol ester in the presence of an inorganic salt such as potassium acetate under the catalysis of a transition metal species such as  
*trans*-dibromobis(triphenylphosphine)palladium (II) or  
[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in a solvent such as  
10 N,N-dimethylformamide at temperatures ranging from 50 to 100°C for a length of time ranging from 1 to 48 hours. The boronate of formula XI may be isolated by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The resulting product can be further purified by  
15 chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.


Alternatively, the boronate of formula XI can be used as generated *in situ* in the reaction medium without isolation, and reacted with a small excess of an appropriately substituted aryl or heteroaryl halide of formula XII under the catalysis  
20 of a transition metal species such as  
[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) in an appropriate solvent or solvent mixture, preferably a 1:1 mixture of toluene and ethanol in the presence of an excess of an aqueous solution of an alkali base such as sodium carbonate, at an appropriate temperature, preferably 50 to 100°C for an appropriate  
25 length of time ranging from 0.5 to 48 hours.

The reaction product of formula I is then isolated in crude form by precipitation with water and filtration or extraction into an appropriate organic solvent such as diethyl ether, ethyl acetate, or a halogenated hydrocarbon solvent such as chloroform or methylene chloride. The product can be further purified by  
30 chromatography on silica gel, crystallization or prolonged stirring in an appropriate solvent followed by filtration.

**Scheme 2****Scheme 3**

- 5 In a third method used for the synthesis of compounds of formula I of this invention (scheme 3), an intermediate nicotinoyl acrylate of formula III is reacted with an appropriately constructed diaryl or heteroarylaryl amine of formula XIII under the conditions described previously to afford a compound of formula XIV which is cyclized by the action of a strong base such as sodium hydride as described above to
- 10 afford an ester of formula X which is processed into a compound of formula I via hydrolysis and amide formation as described above.

Scheme 3




5

Scheme 4

The diaryl or heteroarylamine intermediates of formula XIII were assembled as indicated in Scheme 4. An appropriately substituted aniline boronic acid of formula XV is coupled with an appropriately substituted aryl or heteroaryl halide of general formula XII under the catalysis of a transition metal species as described above to afford the formula XIII compounds used in Scheme 3.


10

Scheme 4

5

**Scheme 5**


Bromopyridine intermediates substituted at the 2-position by carbon based nucleophiles of formula **XVII**, where  $\text{R}^8$  is selected from  $\text{R}^2$  moieties having a carbon-carbon link to the pyridine, are conveniently prepared as shown in **Scheme 5**. The bromopyridine intermediates are prepared from dihalides of formula **XVI** by treatment with an appropriate solution of a Grignard reagent under the catalysis of a transition metal species such as [1,1'-bis(diphenylphosphino)ferrocene]dichloronickel (**II**) in a solvent such as tetrahydrofuran at a temperature range of  $-10^\circ\text{C}$  to room temperature and the resulting reaction mixture worked up by well known procedures to afford the desired product.

Scheme 5

## Scheme 6

- Halopyridine intermediates of type XVIII where the 2-substituent is an alkoxy group OR<sup>9</sup> are derived from dihalides of formula XVI by displacement with an appropriate alkali alkoxide as outlined in scheme 6. The reaction is effected in a solvent such as N,N-dimethylformamide at a temperature range of 0°C to room temperature and, upon completion of the reaction, the products are isolated and purified following classical procedures.

Scheme 6




## Scheme 7

- Where intermediates of formula XIX or XX in which the 2-substituent is sulfide, sulfoxide or sulfone were required, they were attained as described in Scheme 7. An appropriate dihalopyridine of type XVI is reacted with an appropriate thioalkoxide, usually generated from the corresponding thiol or disulfide through the action of a strong base such as an alkali hydride or n-butyllithium, in a solvent such as N,N-dimethylformamide or diethyl ether at a temperature range of -78°C to room temperature. Upon completion of the reaction the products of formula XIX are isolated and purified following classical procedures. The products thus obtained can be oxidized to the corresponding sulfoxides or sulfones of formula XX through the action of an oxidizing agent such as oxone or an organic peracid. In Scheme 7, R<sup>10</sup> is H or C<sub>1-6</sub>alkyl.



Scheme 7




5

Scheme 8

The preparation of haloacyl pyridine intermediates of formula **XXII**, presented in **Scheme 8**, requires treatment of a halopyridine ester of type **XXI** with a solution of an appropriate Grignard reagent in a solvent such as diethyl ether at a temperature range of 0°C to room temperature. If the reaction is carried out for a longer period or under reflux a halopyridine carbinol of formula **XXIII** is obtained. In **Schemes 8** and **9**, R<sup>7</sup> is C<sub>1-6</sub>alkyl and R<sup>6</sup> is methyl or ethyl.


Scheme 8



15

## Scheme 9

- Scheme 9 outlines an alternative sequence for the synthesis of certain halopyridine carbinols of type XXIII. When 2,5-dibromopyridine is treated with n-butyllithium in toluene at  $-78^{\circ}\text{C}$  followed by addition of an appropriate ketone or aldehyde and subsequent quenching at  $-78^{\circ}\text{C}$ , a carbinol of type XXIII results where the carbinol group occupies the 2-position of the pyridine ring. If the metallation step is performed in diethyl ether, the same process leads to an intermediate of formula XXIII in which the carbinol group occupies the 5-position of the pyridine ring.




15

## Scheme 10

- Scheme 10 demonstrates the methods of synthesis for compounds of formula I in which  $\text{R}^2$  is a substituted phenyl or heteroaryl group. An intermediate compound of Type I where  $\text{R}^2$  is a halogen is reacted with an appropriately substituted boronic acid or boronate ester of formula VII or tributyl stannane of formula IX using one of the methods described above to afford the desired compound.

Scheme 10



Scheme 11

5

Various further transformations on pre-assembled compounds of formula **I** are illustrated in **Scheme 11**. In cases where the Ar group is a pyridine or quinoline group it can be oxidized to the corresponding nitrogen oxide by the action of an appropriate oxidizing agent such as m-chloroperoxybenzoic acid or magnesium monoperoxyphthalate under commonly used conditions. In cases where one or more of the substituents on the Ar group is a ketone it is conveniently transformed into an oxime analog through the action of hydroxylamine in pyridine as solvent. A sulfide substituent is easily oxidized to the corresponding sulfoxide or sulfone derivative by using an appropriate quantity of an oxidant such as oxone or an organic peracid.


15

The transformation of a 2-benzyloxypyridine into the corresponding 2-pyridone was accomplished by treatment with trifluoroacetic acid in a solvent such as methylene chloride at room temperature or under slight warming. The removal of a tert-butyloxycarbonyl protecting group from a piperazine ring is effected by reaction with trifluoroacetic acid in a solvent such as 1,2-dichloroethane at reflux temperature. In examples where a substituent on Ar is a hydroxymethyl group it can be derivatized to the analogous halomethyl moiety using a tetrahalomethane in the presence of a trisubstituted phosphine such as triphenylphosphine or diphos in a solvent such as

20

methylene chloride. The halide can be displaced by an appropriate sulfinic acid sodium salt to afford the alkyl or arylsulfonylmethyl analog.


Scheme 11



5

10





I

EX.	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>6</sup>
1	H	i-pr	3-C(O)Me	H	H	H
2	H		3-C(O)Me	H	H	H
3	H	i-pr	4-n-propyl	H	H	H
4	H	i-pr	4-C(O)Me	H	H	H
5	H	i-pr	2-Me	H	H	H
6	Me	i-pr	4-C(O)Me	H	H	H
9	H	t-bu	4-C(O)Me	H	H	H
11	H	i-pr	4-	H	H	H
16	H	c-pr	4-CH <sub>2</sub> OH	H	H	H
18	H	c-pr	4-SEt	H	H	H
20	H	c-pr	4-SO <sub>2</sub> NH <sub>2</sub>	H	H	H
21	H	i-pr	3-OEt	H	H	H
22	H	i-pr	4-SMe	H	H	H
23	H	i-pr	3-C(O)Me	4-OH	H	H
49	H	i-pr	4-SO <sub>2</sub> Me	H	H	H
52	H	c-pr	4-SO <sub>2</sub> Et	H	H	H





EX.	R	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>6</sup>
53	H	c-pr	4-S(O)Et	H	H	H
54	H	i-pr	4-C(=NOH) Me	H	H	H
55	H	i-pr	4- 	H	H	H
56	H	c-pr	4-CH <sub>2</sub> SO <sub>2</sub> Me	H	H	H

Table 2

5




EX.	R	R <sup>1</sup>	$\pi$ - Position	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>6</sup>	n
7	H	i-pr	3	H	H	H	H	0
10	H		3	H	H	H	H	0
14	H	c-pr	3	H	H	H	H	0
15	H	i-pr	3	5-SMe	H	H	H	0
17	H	c-pr	4	H	H	H	H	0
24	H	i-pr	3	5-COOEt	H	H	H	0

EX.	R	R <sup>1</sup>	$\pi$ - Position	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>6</sup>	n
25	H	i-pr	3	5-CMe <sub>2</sub> OH	H	H	H	0
26	H	i-pr	3	6-CH <sub>2</sub> CHMe <sub>2</sub>	H	H	H	0
27	H	i-pr	3	5-C(O)Me	H	H	H	0
28	H	i-pr	3	6-Me	H	H	H	0
30	H	H	3	6-CMe <sub>2</sub> OH	H	H	H	1
32	H	c-pr	3	5-SO <sub>2</sub> Me	H	H	H	0
33	H	c-pr	2	4-CMe <sub>2</sub> OH	H	H	H	1
34	H	c-pr	2	5-CMe <sub>2</sub> OH	H	H	H	0
35	H	c-pr	4	3-CMe <sub>2</sub> OH	H	H	H	0
36	H	c-pr	4	3-CMe <sub>2</sub> OH	H	H	H	1
37	H	c-pr	3	6-SO <sub>2</sub> I-pr	H	H	H	0
38	H	c-pr	3	6-OMe	H	H	H	0
39	H	c-pr	3	6-Me	H	H	H	0
40	H	c-pr	3	6-OCH <sub>2</sub> CF <sub>3</sub>	H	H	H	0
41	H	c-pr	3	5-Br	H	H	H	0
42	H	c-pr	3	6-OCH <sub>2</sub> Ph	H	H	H	0
43	H	c-pr	3	6-C(c-pr) <sub>2</sub> OH	H	H	H	0
44	H	c-pr	2	5-CMe <sub>2</sub> OH	H	H	H	1
45	H	c-pr	3	6-CMe <sub>2</sub> OH	H	H	H	0
46	H	i-butyl	3	6-CMe <sub>2</sub> OH	H	H	H	0
47	H	c-pr	3	6-CMe <sub>2</sub> OH	H	H	5-Br	0
48	H	c-pr	2	6-CMe <sub>2</sub> OH	H	H	H	0
50	H	c-pr	3	6-SO <sub>2</sub> Me	H	H	H	0
51	H	i-pr	3	5-SO <sub>2</sub> Me	H	H	H	0
59	H	i-pr	3	H	H	H	H	1
60	H		3	H	H	H	H	1
61	H	i-pr	3	5-COOEt	H	H	H	1
62	H	i-pr	3	5-CMe <sub>2</sub> OH	H	H	H	1




EX.	R	R <sup>1</sup>	$\pi$ - Position	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>6</sup>	n
63	H	i-pr	3	6-CH <sub>2</sub> CHMe <sub>2</sub>	H	H	H	1
64	H	i-pr	3	6-Me	H	H	H	1
65	H	c-pr	3	H	H	H	H	1
66	H	c-pr	3	6-CMe <sub>2</sub> OH	H	H	H	1
67	H	c-pr	4	H	H	H	H	1
68	H	c-pr	3	5-Br	H	H	H	1
73	H	i-butyl	3	6-CMe <sub>2</sub> OH	H	H	H	1
74	H	c-pr	3	6-Me	H	H	H	1
75	H	c-pr	3	6-SO <sub>2</sub> Me	H	H	H	1
76	H	c-pr	3	6-CMe <sub>2</sub> OH	H	H	5-Br	1
77	H	c-pr	3	6-CMe(CH <sub>2</sub> OH)OH	H	H	H	1


Table 3







5

Example	R	R <sup>1</sup>	Ar	R <sup>4</sup>	R <sup>6</sup>
8	H	i-pr	indol-5-yl	H	H
12	H	i-pr	quinolin-3-yl	H	H
13	H	i-pr	pyrimidin-5-yl	H	H
19	H	c-pr	3-thienyl	H	H
29	H	c-pr	1-oxidopyrimidin-5-yl	H	H


Example	R	R <sup>1</sup>	Ar	R <sup>4</sup>	R <sup>6</sup>
57	H	c-pr		H	H
72	H	i-pr	1-oxidoquinolin-3-yl	H	H

**Table 4**











5










Example	R	R <sup>1</sup>	Ar	Ar <sup>1</sup>	R <sup>4</sup>	R <sup>6</sup>
31	H	i-pr	Ph	4-(pyridin-3-yl)	H	H
58	H	c-pr	Pyridin-3-yl		H	H
69	H	c-pr	Pyridin-3-yl		H	H
70	H	c-pr	1-oxidopyridin-3-yl		H	H
71	H	c-pr	1-oxidopyridin-3-yl		H	H



**Table 5**



R <sup>1</sup>	R <sup>4</sup>	R <sup>6</sup>	Ar
c-pr	H	H	
c-pr	H	H	
i-pr	7-Me	H	
i-pr	H	5-Me	
c-pr	H	H	

R <sup>1</sup>	R <sup>4</sup>	R <sup>6</sup>	Ar
c-pr	H	H	
c-pr	H	H	
c-Bu	H	6-F	
	6-F	H	
	H	H	
	5-F	4-F	
i-pr	H	H	


R <sup>1</sup>	R <sup>4</sup>	R <sup>6</sup>	Ar
i-pr	H	H	
i-pr	H	H	
c-pr	H	H	
	H	H	
c-pr	H	H	
c-Bu	H	H	
	H	H	

R <sup>1</sup>	R <sup>4</sup>	R <sup>6</sup>	Ar
	H	H	

## EXAMPLE 1

5

**N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: Ethyl 3-(3-bromoanilino)-2-(2-chloronicotinoyl) acrylate.**

10

A mixture of ethyl 2-chloronicotinoyl acetate (41.1g, 180.5mmol), triethyl orthoformate (40.12g, 271mmol) and acetic anhydride (92.05g, 902.5mmol) was heated at 130°C for 2.5 hours. The volatile components were distilled off and the residue was co-evaporated twice with xylene. The oily residue was dissolved in methylene chloride (250mL) and 3-bromoaniline (37.25g, 216.6mmol) was added slowly. The resulting solution was stirred at room temperature for 18 hours, and the solvent evaporated away. The resulting crude compound was used as such in the next step.

Step 2: Ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate.

The crude compound from **Step 1** was dissolved in tetrahydrofuran (500mL), the solution was cooled to 0°C, and sodium hydride (as a 60% dispersion in oil, 9.4g, 235mmol) was added in portions. After stirring at 0° for 1 hour, the mixture was allowed to warm up to room temperature. After 2 hours, water (400mL) was added to the suspension and the insoluble solid was filtered and washed copiously with water. When dry, the solid was stirred in ether (150mL) at room temperature for 24 hours and filtered to afford the title compound as a cream-colored solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.32 (t, 3H), 4.29 (q, 2H), 7.54-7.63 (m, 2H), 7.69 (dd, 1H), 7.78 (dd, 1H), 7.93 (s, 1H), 8.66-8.71 (m, 3H).

Step 3: 1-(3-Bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid

A suspension of ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from **Step 2** (52.5g, 140.7mmol) in a mixture of tetrahydrofuran (400mL), methanol (400mL) and 1N aqueous sodium hydroxide (280mL) was heated at ca 50°C with stirring for 20 minutes. After cooling, the mixture was diluted with water (300mL) and 1N aqueous HCl (325mL) was added. After stirring for 45 minutes, the precipitate was filtered, washed well with water and dried to afford the title acid as a cream-colored solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 7.65 (t, 1H), 7.76 (m, 2H), 7.84 (d, 1H), 7.99 (s, 1H), 8.87 (m, 2H), 9.01 (s, 1H).

Step 4: N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.

To a suspension of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from **Step 3** (26.3g, 76mmol) and triethylamine (23.2g, 230mmol) in tetrahydrofuran (1000mL) at 0°C was added isobutyl chloroformate (18.85g, 138mmol). After stirring at 0°C for 2 hours, isopropylamine (23g, 390mmol) was added and the mixture was allowed to warm up to room temperature and stirred

overnight. The mixture was then partitioned between ethyl acetate and water, the organic phase was dried and evaporated to a solid which was stirred in ether at room temperature for 3 hours and filtered to afford the N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide as a white solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.25 (d, 6H), 4.17 (m, 1H), 7.59-7.63 (m, 2H), 7.70 (d, 1H), 7.80 (d, 1H), 7.94 (s, 1H), 8.73 (m, 1H), 8.78 (d, 1H), 8.85 (s, 1H), 9.61 (br, NH).

**Step 5: N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


A mixture of N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 4, 3-acetylphenylboronic acid (1.2eq.), *trans*-dibromobis(triphenylphosphine)palladium (II) (0.05eq.), toluene (6mL/mmol), ethanol (2mL/mmol) and 2M aqueous sodium carbonate (8eq.) was refluxed for 1 hour under a nitrogen atmosphere. The mixture was diluted with ethyl acetate and the organic phase was washed with water and brine, dried and evaporated. The crude product was chromatographed on silica gel eluting with a gradient of 20-40% ether in methylene chloride to afford the N-Isopropyl-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide product as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 2.65 (s, 3H), 4.28 (m, 1H), 7.47 (m, 2H), 7.55 (t, 1H), 7.65 (m, 2H), 7.80 (m, 2H), 7.95 (dd, 1H), 8.19 (brs, 1H), 8.70 (dd, 1H), 8.81 (dd, 1H), 9.05 (s, 1H), 9.65 (br, NH).

**EXAMPLE 2**

**N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





**Step 1: Anion of 4-amino-3,5-dichloropyridine.**

A suspension of sodium hydride as 60% dispersion in oil (360mg, 9mmol) in tetrahydrofuran (15mL) was cooled to 0°C. A solution of 4-amino-3,5-dichloropyridine (978mg, 6mmol) in tetrahydrofuran (15mL) was added slowly. The resulting mixture was kept at 0° for 2.5 hours.

**Step 2: Acid chloride of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid.**

A suspension of 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 3 of Example 1 (690mg, 2mmol) in tetrahydrofuran (12mL) was cooled to 0°C, and oxalyl chloride (381mg, 3mmol) was added, followed by 2 drops of N,N-dimethylformamide. The resulting mixture was then stirred at room temperature for 1 hour then refluxed for 45 minutes and cooled to room temperature.

**Step 3: N-(2,6-Dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.**

The mixture from Step 2 above, as a brown suspension, was added via syringe to the cold suspension of Step 1. The resulting mixture was stirred at room temperature for 18 hours, quenched with aqueous saturated ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from evaporation of the organic phase was triturated with ether (50mL) and filtered,

affording the N-(2,6-Dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide as a beige solid.

$^1\text{H}$  NMR (Acetone- $d_6$ )  $\delta$  7.61-7.70 (m, 2H), 7.76 (d, 1H), 7.81 (d, 1H), 8.00 (s, 1H), 8.62 (s, 2H), 8.80 (br s, 1H), 8.86 (d, 1H), 8.99 (s, 1H), 12.1 (br, NH).

5

**Step 4: N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.**


Following the procedure of **Step 5 of Example 1**, but substituting N-(2,6-dichloropyridin-4-yl)-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 3 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-(2,6-Dichloropyridin-4-yl)-1-[3-(3-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.65 (s, 3H), 7.47 (d, 1H), 7.50-7.60 (m, 2H), 7.70 (m, 2H), 7.82 (d, 2H), 7.98 (d, 1H), 8.20 (s, 1H), 8.55 (s, 2H), 8.75 (brs, 1H), 8.92 (dd, 1H), 9.14 (s, 1H), 12.08 (br, NH).

20

**EXAMPLE 3**

**N-Isopropyl-1-[3-(4-n-propylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5** of **Example 1**, but substituting 4-n-propylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a white solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.93 (t, 3H), 1.24 (d, 6H), 1.65 (m, 2H), 2.62 (t, 2H), 4.18 (m, 1H), 7.31 (d, 2H), 7.58-7.61 (m, 2H), 7.68-7.72 (m, 3H), 7.87 (d, 1H), 7.95 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.66 (br, NH).

#### EXAMPLE 4

**N-Isopropyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5** of **Example 1**, but substituting 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.25 (d, 6H), 2.61 (s, 3H), 4.17 (m, 1H), 7.59 (m, 1H), 7.70 (d, 1H), 7.76 (t, 1H), 7.92 (d, 2H), 7.97 (d, 1H), 8.07-8.10 (m, 3H), 8.72 (brs, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.65 (br, NH).

#### EXAMPLE 5

**N-Isopropyl-1-[3-(2-methylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5 of Example 1**, but substituting 2-methylphenylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

$^1\text{H}$  NMR (Acetone- $d_6$ )  $\delta$  1.24 (d, 6H), 2.35 (s, 3H), 4.17 (m, 1H), 7.27-7.34 (m, 4H), 7.56-7.60 (m, 2H), 7.65 (m, 2H), 7.70 (t, 1H), 8.74 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 9.64 (br, NH).

**EXAMPLE 6**

**N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: N-Isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Example 1, Step 4**, but substituting N-isopropyl-N-methylamine for isopropylamine the N-Isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained in as a yellow solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) (Appears as two rotamers of the amide) δ 1.18 (m, 6H), 2.85 (s, 3H), 4.05 (m, 0.5H), 4.84 (m, 0.5H), 7.49-7.64 (m, 3H), 7.72 (d, 1H), 7.86 (s, 1H), 8.14 (s, 1H), 8.65 (d, 2H).


**Step 2: N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 5 of Example 1**, but substituting N-isopropyl-N-methyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the N-Isopropyl-N-methyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) (Appears as two rotomers of the amide) δ 1.23 (m, 6H), 2.62 (s, 3H), 4.00 (m, 0.5H), 4.92 (m, 0.5H), 7.38-7.55 (m, 2H), 7.63-7.77 (m, 5H), 8.03 (d, 2H), 8.14 (s, 0.5H), 8.21 (s, 0.5H), 8.65 (m, 1H), 8.75-8.80 (m, 1H).

**EXAMPLE 7**

**N-Isopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5 of Example 1**, but substituting pyridine-3-boronic acid 1,3-propanediol cyclic ester for 3-acetylphenylboronic acid and [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) the title compound was obtained as a beige solid.

$^1\text{H}$  NMR (Acetone- $d_6$ )  $\delta$  1.24 (d, 6H), 4.17 (m, 1H), 7.48 (m, 1H), 7.60 (m, 1H), 7.71 (dd, 1H), 7.78 (t, 1H), 7.95 (dd, 1H), 8.05 (brs, 1H), 8.15 (m, 1H), 8.60 (m, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.92 (s, 1H), 8.99 (brs, 1H), 9.65 (br, NH).

## EXAMPLE 8

**N-Isopropyl-1-[3-(indol-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5 of Example 1**, but substituting 5-indolylboronic acid for 3-acetylphenylboronic acid the title compound was obtained as an off-white solid.

<sup>1</sup>H NMR ((DMSO-d<sub>6</sub>) δ 1.20 (d, 6H), 4.10 (m, 1H), 6.47 (s, 1H), 7.38 (brs, 1H), 7.46-7.52 (m, 3H), 7.59-7.66 (m, 2H), 7.87-7.93 (m, 3H), 8.72-8.81 (m, 3H), 9.67 (br, NH), 11.2 (br, NH).

### EXAMPLE 9

10

**N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: N-tert-Butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

15

Following the procedure of **Example 1, Step 4**, but substituting tert-butylamine for isopropylamine the N-tert-Butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained as a yellow solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.44 (s, 9H), 7.58-7.62 (m, 2H), 7.70 (dd, 1H), 7.78 (dd, 1H), 7.93 (br s, 1H), 8.72 (m, 1H), 8.77 (dd, 1H), 8.81 (s, 1H), 9.73 (br, NH).

20


**Step 2: N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 5** of **Example 1**, but substituting N-tert butyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-acetylphenylboronic acid for 3-acetylphenylboronic acid the N-tert-Butyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained in 93% yield as a white solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.45 (s, 9H), 2.61 (s, 3H), 7.59 (m, 1H), 7.69-7.72 (m, 1H), 7.77 (t, 1H), 7.92-7.99 (m, 3H), 8.07-8.11 (m, 3H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.91 (s, 1H), 9.79 (br, NH).

**EXAMPLE 10**

**N-(2,6-Dichloropyridin-4-yl)-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



Following the procedure of **Step 4** of **Example 2** but substituting [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) and pyridine-3-boronic acid 1,3-propanediol cyclic ester for 3-acetylphenylboronic acid the title compound was obtained as a glassy solid.




<sup>1</sup>H NMR (Acetone-*d*<sub>6</sub>) δ 7.48 (m, 1H), 7.68 (m, 1H), 7.77-7.82 (m, 2H), 7.98 (m, 1H), 8.12-8.17 (m, 2H), 8.61 (m, 1H), 8.62 (s, 2H), 8.80 (m, 1H), 8.88 (dd, 1H), 8.99 (brs, 1H), 9.06 (s, 1H), 12.2 (br, NH).

5

### EXAMPLE 11

**N-Isopropyl-1-{3-[ 4-(4-*tert*butyloxycarbonylpiperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



10

**Step 1: 4-*tert*-Butyloxycarbonyl-1-(3-bromophenyl)piperazine**

To a suspension of 1-(4-bromophenyl)piperazine hydrochloride (103.15g, 371.59mmol) in acetonitrile (1.5 L) at 0°C under a nitrogen atmosphere was added a catalytic amount of 4-dimethylaminopyridine (4.54g, 37.159mmol) followed by triethylamine (155mL, 1114.77mmol) and di-*tert*-butyl dicarbonate (121.65g, 557.385mmol, dissolved in a minimum amount of acetonitrile) and the resulting reaction mixture was warmed to room temperature and stirred for 5.5 hours. The reaction mixture was filtered, ethyl acetate was added and the organic phase was washed with 10 % aqueous citric acid, water (2X) and brine, then dried and

evaporated to afford the crude 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine product which was used as such in the next step.

**Step 2: 3-(4-tert-Butyloxycarbonylpiperazin-1-yl)phenylboronic acid**

5                   To the 4-tert-Butyloxycarbonyl-1-(3-bromophenyl)piperazine from  
Step 1 (118.30g, 346.9mmol) in tetrahydrofuran /toluene (1/1, 1.5 L) at  $-78^{\circ}\text{C}$  under  
nitrogen was added n-butyllithium (2.5M, 160mL, 398.9mmol) dropwise and the  
resulting reaction mixture was stirred at  $-78^{\circ}\text{C}$  for 20 minutes. Triisopropyl borate  
(96.1mL, 416.3mmol) was added dropwise and the reaction was warmed to  $0^{\circ}\text{C}$  and  
10                   stirred for 2 hours. Aqueous saturated ammonium chloride (400mL), water (100mL)  
and 1 equivalent of  $\text{H}_3\text{PO}_4$  (20mL) were added and the mixture was stirred for 15  
minutes and then concentrated to a volume of approximately 200mL (at which stage  
the mixture became bluish and a precipitate formed). The mixture was slowly diluted  
with heptane (800mL) and the resulting suspension was stirred overnight. The  
15                   suspension was filtered, the solid was washed with heptane and dried to afford the  
title boronic acid.


**Step 3: N-Isopropyl-1-[3-[4-(4-tertbutyloxycarbonylpiperazin-1-yl)phenyl]-  
phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

20                   Following the procedure of Step 5 of Example 1 but substituting  
[1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-  
dibromobis(triphenylphosphine)palladium (II) and the boronic acid from Step 2 above  
for 3-acetylphenylboronic acid the 4-tert-Butyloxycarbonyl-1-(3-  
bromophenyl)piperazine compound was obtained as a solid.

25                    $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.30 (d, 6H), 1.49 (s, 9H), 3.18 (m, 4H), 3.58 $\checkmark$  (m,  
4H), 4.29 (m, 1H), 6.98 (d, 2H), 7.32 (d, 1H), 7.45 (m, 1H), 7.53 (d, 2H), 7.55-7.62  
(m, 2H), 7.72 (d, 1H), 8.70 (m, 1H), 8.82 (d, 1H), 9.07 (s, 1H), 9.68 (br, NH).

30                   **EXAMPLE 12**

**N-Isopropyl-1-[3-(quinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-  
carboxamide**




Following the procedure of **Step 5 of Example 1**, but substituting 3-quinolineboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 4.29 (m, 1H), 7.49 (m, 2H), 7.61 (t, 1H), 7.70-7.78 (m, 3H), 7.86-7.92 (m, 2H), 8.14 (d, 1H), 8.36 (s, 1H), 8.71 (m, 1H), 8.84 (dd, 1H), 9.10 (s, 1H), 9.19 (s, 1H), 9.67 (br, NH).

10

### EXAMPLE 13

**N-Isopropyl-1-[3-(pyrimidin-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide**




15

Following the procedure of **Step 5 of Example 1**, but substituting 5-pyrimidineboronic acid for 3-acetylboronic acid the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.28 (d, 6H), 4.27 (m, 1H), 7.48 (dd, 1H), 7.52 (m, 1H), 7.65 (s, 1H), 7.74 (m, 2H), 8.68 (m, 1H), 8.72 (d, 1H), 8.98 (s, 2H) 9.03 (s, 1H), 9.22 (s, 1H), 9.62 (br, NH).

5

**EXAMPLE 14****N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

10

**Step 1: N-Cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Example 1, Step 4**, but substituting cyclopropylamine for isopropylamine the N-Cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide was obtained as a fluffy white solid.

15

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.59 (m, 2H), 0.80 (m, 2H), 2.96 (m, 1H), 7.59-7.68 (m, 2H), 7.72 (dd, 1H), 7.82 (dd, 1H), 7.97 (s, 1H), 8.72-8.81 (m, 2H), 8.89 (s, 1H), 9.70 (br, NH).

**Step 2: N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Example 7** but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-


25

dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a cream-coloured solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 7.52 (m, 1H), 7.63-7.69 (m, 2H), 7.74 (t, 1H), 7.97 (d, 1H), 8.07 (brs, 1H), 8.17 (d, 1H), 8.61 (m, 1H), 8.73 (dd, 1H), 8.79 (m, 1H), 8.85 (s, 1H), 8.99 (brs, 1H), 9.74 (br, NH).

### EXAMPLE 15

#### N-Isopropyl-1-[3-(5-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




Following the procedure of Step 5 of Example 1 but substituting 5-methylthiopyridine-3-boronic acid for 3-acetylphenylboronic acid and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) for *trans*-dibromobis(triphenylphosphine)palladium (II) the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.33 (d, 6H) 2.60 (s, 3H), 4.33 (m, 1H), 7.48-7.54 (m, 2H), 7.66 (m, 1H), 7.73 (t, 1H), 7.78-7.81 (m, 2H), 8.55 (s, 1H), 8.66 (s, 1H), 8.74 (m, 1H), 8.87 (d, 1H), 9.09 (s, 1H), 9.69 (br, NH).

### EXAMPLE 16

**N-Cyclopropyl-1-[3-(4-hydroxymethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2 of Example 14** but substituting 4-hydroxymethylphenyl boronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.71 (m, 2H), 0.89 (m, 2H), 1.88 (t, 1H), 3.03 (m, 1H), 4.78 (d, 2H), 7.43 (d, 1H), 7.46-7.52 (m, 3H), 7.61-7.69 (m, 4H), 7.80 (d, 1H), 8.73 (m, 1H), 8.83 (dd, 1H), 9.10 (s, 1H), 9.82 (br, NH).

**EXAMPLE 17**

**N-Cyclopropyl-1-[3-(pyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 5** of **Example 1** but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide and 4-pyridineboronic acid for 3-acetylphenylboronic acid the title compound was obtained as a white solid.

$^1\text{H}$  NMR ((DMSO- $d_6$ )  $\delta$  0.57 (m, 2H), 0.77 (m, 2H), 2.90 (m, 1H), 7.64 (m, 1H), 7.72-7.89 (m, 4H), 8.03 (d, 1H), 8.13 (s, 1H), 8.66-8.78 (m, 4H), 8.84 (s, 1H), 9.72 (br, NH).

### EXAMPLE 18

**N-Cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2** of **Example 14** but substituting 4-ethylthiobenzeneboronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.72 (m, 2H), 0.90 (m, 2H), 1.48 (t, 3H), 3.03 (m, 3H), 7.42 (d, 3H), 7.50 (m, 1H), 7.57 (d, 2H), 7.64 (s, 1H), 7.68 (t, 1H), 7.78 (d, 1H), 8.75 (m, 1H), 8.85 (d, 1H), 9.10 (s, 1H), 9.83 (br, NH).

### EXAMPLE 19

**N-Cyclopropyl-1-[3-(3-thienyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2** of **Example 14** but substituting 3-thiopheneboronic acid for pyridine-3-boronic acid 1,3-propanediol cyclic ester the title compound was obtained as a white solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.60 (m, 2H), 0.79 (m, 2H), 2.96 (m, 1H), 7.57-7.72 (m, 5H), 7.92-7.98 (m, 2H), 8.05 (s, 1H), 8.74 (s, 1H), 8.78 (d, 1H), 8.93 (s, 1H), 9.74 (br, NH).

**EXAMPLE 20**

**N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





**Step 1: 4-Sulfamoylbenzeneboronic acid pinacol ester**

A mixture of 4-bromobenzenesulfonamide, diboron pinacol ester (1.1eq), potassium acetate (3.5eq) and 1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-dimethylformamide (4ml/mmol) was heated at 85°C for 18 hours. After quenching with saturated aqueous ammonium chloride solution the mixture was partitioned between ethyl acetate and water and the product from the organic phase was chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and hexane to afford the 4-Sulfamoylbenzeneboronic acid pinacol ester as a solid.


**Step 2: N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

A mixture of N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, boronate from Step 1 (1.2eq), palladium acetate (0.1eq), triphenylphosphine (0.35eq) and 2M aqueous sodium carbonate (3.5eq) in n-propanol (10ml/mmol) was stirred at 85°C for 1 hour. After cooling, the mixture was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water, and the product from the organic phase was chromatographed on silica gel eluting with a 1:5:4 mixture of ethanol, ethyl acetate and methylene chloride to afford the N-Cyclopropyl-1-[3-(4-sulfamoylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.62 (m, 2H), 0.82 (m, 2H), 2.98 (m, 1H), 6.66 (br, NH<sub>2</sub>), 7.64 (m, 1H), 7.74 (m, 1H), 7.80 (t, 1H), 7.97-8.05 (m, 5H), 8.10 (m, 1H), 8.76 (m, 1H), 8.81 (dd, 1H), 8.97 (s, 1H), 9.77 (br, NH).

**EXAMPLE 21**

**N-Isopropyl-1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: Ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate**

Following the procedure of **Step 5 of Example 1**, but substituting ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from **Step 2 of Example 1** for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, and 3-ethoxybenzeneboronic acid for 3-acetylbenzeneboronic acid, the Ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate compound was obtained as a solid.

**Step 2: 1-[3-(3-Ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid**

Following the procedure of **Step 3 of Example 1** but substituting ethyl 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from step 1 for ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate the 1-[3-(3-Ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained and used without purification in the next step.

**Step 3: N-Isopropyl- 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


A mixture of 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from **Step 2** and thionyl chloride (4eq) in tetrahydrofuran (10ml/mmol) was refluxed for 45 minutes, then evaporated. The residue was dissolved in the same volume of tetrahydrofuran, isopropylamine (5eq) was added and the mixture was stirred at room temperature for 18 hours. After

quenching with saturated aqueous ammonium chloride solution, the resulting mixture was partitioned between ethyl acetate and water, and the product from the organic phase was chromatographed on silica gel eluting with 10% ether in methylene chloride to afford the N-Isopropyl- 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.29 (d, 6H), 1.42 (t, 3H), 4.08 (q, 2H), 4.28 (m, 1H), 6.91 (d, 1H), 7.12 (s, 1H), 7.18 (d, 1H), 7.34 (t, 1H), 7.40 (d, 1H), 7.46 (m, 1H), 7.60-7.65 (m, 2H), 7.75 (d, 1H), 8.71 (brs, 1H), 8.82 (dd, 1H), 9.08 (s, 1H), 9.70 (br, NH).

## EXAMPLE 22

N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



### Step 1: Ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate

Following the procedure of **Step 5** of **Example 1**, but substituting ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from step 2 of example 1 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, and 4-methylthiobenzeneboronic acid for 3-acetylbenzeneboronic acid, the Ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate compound was obtained as a solid.

**Step 2: 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid**

- Following the procedure of **Step 3 of Example 1** but substituting ethyl 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate from **Step 1** for ethyl 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylate the 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a solid.


**Step 3: N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

- Following the procedure of **Step 3 of Example 21** but substituting 1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid for 1-[3-(3-ethoxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid the N-Isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

- <sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.24 (d, 6H), 2.52 (s, 3H), 4.18 (m, 1H), 7.37 (d, 2H), 7.58-7.62 (m, 2H), 7.69-7.73 (m, 3H), 7.87 (d, 1H), 7.96 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.91 (s, 1H), 9.65 (br, NH).

**EXAMPLE 23**

- N-Isopropyl-1-[3-(3-acetyl-4-hydroxyphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




A mixture of 5'-bromo-2'-hydroxyacetophenone, diboron pinacol ester (1.25eq), potassium acetate (3eq) and [1,1'-bis  
(diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-  
5 dimethylformamide (10ml/mmol) was stirred at 80°C for 3 hours and cooled down. A solution of N-Isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-  
carboxamide from **Example 1, Step 4** (0.75eq) in N,N-dimethylformamide  
(7ml/mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq)  
and 2M aqueous sodium carbonate (8.5eq) were added and the resulting mixture was  
10 stirred at 80°C for 2.5 hours. The cooled mixture was partitioned between ethyl  
acetate and water and the product from the organic phase was chromatographed on  
silica gel eluting with 60% ethyl acetate in hexane to afford the title compound as a  
light yellow solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.24 (d, 6H), 2.75 (s, 3H), 4.19 (m, 1H),  
15 7.06 (d, 1H), 7.59-7.63 (m, 2H), 7.72 (t, 1H), 7.92 (d, 1H), 7.97 (d, 1H), 8.02 (s, 1H),  
8.33 (s, 1H), 8.73 (m, 1H), 8.78 (dd, 1H), 8.90 (s, 1H), 9.65 (br, NH).

#### EXAMPLE 24

20

**N-Isopropyl-1-[3-(5-carboethoxyphenyl)phenyl]-1,4-  
dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Example 23** but substituting N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for 5'-bromo-2'-hydroxyacetophenone and ethyl 5-bromonicotinate for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide the title compound was obtained as a beige solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.29 (d, 6H), 1.40 (t, 3H), 4.28 (m, 1H), 4.42 (q, 2H), 7.45-7.51 (m, 2H), 7.68 (s, 1H), 7.71 (t, 1H), 7.80 (d, 1H), 8.49 (s, 1H), 8.59 (m, 1H), 8.82 (d, 1H), 9.03 (s, 1H), 9.07 (s, 1H), 9.23 (s, 1H), 9.64 (br, NH).

#### EXAMPLE 25

**N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 3-Bromo-5-(1-hydroxy-1-methylethyl)pyridine**

To a solution of ethyl 5-bromonicotinate (1.02g, 4.4mmol) in diethyl ether (15ml) at  $-30^{\circ}\text{C}$  was added a 3M solution of methyl magnesium bromide (4ml, 12mmol) in ether. The resulting slurry was then refluxed for 2 hours then cooled and quenched with an excess of 0.5M aqueous monobasic sodium phosphate and partitioned between ether and water. The product from the organic phase was chromatographed on silica gel eluting with a 2:1:2 mixture of ether, pentane and ammonia-saturated methylene chloride to afford the 3-Bromo-5-(1-hydroxy-1-methylethyl)pyridine compound as a yellow oil.


**Step 2: N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.**

Following the procedure of **Example 24**, but substituting the 3-bromo-5-(1-hydroxy-1-methylethyl)pyridine from **Step 1** for ethyl 5-bromonicotinate, the title compound was obtained as a yellow foam.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.28 (d, 6H), 1.62 (s, 6H), 2.52 (brs, 1H), 4.25 (m, 1H), 7.41-7.48 (m, 2H), 7.60-7.68 (m, 2H), 7.75 (d, 1H), 8.05 (s, 1H), 8.67-8.71 (m, 3H), 8.80 (dd, 1H), 9.03 (s, 1H), 9.66 (br, NH).

**EXAMPLE 26**

**N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 5-Bromo-2-(2-methylpropyl)pyridine**

To a solution of 2,5-dibromopyridine (4.5g, 19mmol) in tetrahydrofuran (50ml) was added [1,1'-bis (diphenylphosphino)ferrocene]dichloronickel (II) (103mg, 0.19mmol) and the resulting mixture was cooled to  $-10^{\circ}\text{C}$ . A 2M solution of isobutylmagnesium bromide in ether (12.4ml, 24.7mmol) was added slowly and the mixture was stirred at  $-10$  to  $10^{\circ}\text{C}$  for 3.5 hours. After quenching with saturated aqueous ammonium chloride solution, the mixture was partitioned between ether and water and the product from the organic phase was chromatographed on silica gel eluting with 10% ether in pentane to afford the 5-Bromo-2-(2-methylpropyl)pyridine compound as a volatile oil.

**Step 2: N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


Following the procedure of **Example 24** but substituting 5-bromo-2-(2-methylpropyl)pyridine from **Step 1** for ethyl 5-bromonicotinate the N-Isopropyl-1-{3-[6-(2-methylpropyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.92 (d, 6H), 1.28 (d, 6H), 2.10 (m, 1H), 2.69 (d, 2H), 4.28 (m, 1H), 7.19 (d, 1H), 7.40-7.47 (m, 2H), 7.60 (s, 1H), 7.64 (t, 1H), 7.73 (d, 1H), 7.79 (dd, 1H), 8.68 (m, 1H), 8.77-8.83 (m, 2H), 9.05 (s, 1H), 9.66 (br, NH).

**EXAMPLE 27**

**N-Isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





### Step 1: 3-Acetyl-5-bromopyridine

To a solution of ethyl 5-bromonicotinate (3.9g, 16.9mmol) in ether (50ml) at 0°C was added a 3M solution of methylmagnesium bromide (16.9ml, 50.8mmol). The resulting thick slurry was warmed slowly to room temperature and after 1.5 hours it was poured slowly into an excess of 1M aqueous monobasic sodium phosphate. The mixture was partitioned between ether and water and the product from the organic phase was chromatographed on silica gel, eluting with a 1:1:2 mixture of ether, pentane and ammonia-saturated methylene chloride to afford the 3-acetyl-5-bromopyridine compound. This preparation also afforded 3-bromo-5-(1-hydroxy-1-methylethyl)pyridine described in **Example 25**.

### Step 2: N-Isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide


Following the procedure of **Example 23** but substituting 3-acetyl-5-bromopyridine from **Step 1** for ethyl 5-bromonicotinate the N-isopropyl-1-[3-(5-acetylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 2.69 (s, 3H), 4.28 (m, 1H), 7.48 (dd, 1H), 7.51 (d, 1H), 7.69 (s, 1H), 7.72 (t, 1H), 7.80 (d, 1H), 8.42 (s, 1H), 8.69 (m, 1H), 8.82 (d, 1H), 9.05 (s, 2H) 9.17 (s, 1H), 9.63 (br, NH).

### **EXAMPLE 28**

25

**N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 5-Bromo-2-methylpyridine**

5                      Following the procedure of **Step 1** of **Example 26** but substituting methylmagnesium chloride for isobutylmagnesium bromide the 5-bromo-2-methylpyridine compound was obtained as a solid.

**Step 2: N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


10                      Following the procedure of **Example 24** but substituting 5-bromo-2-methylpyridine from **Step 1** for ethyl 5-bromonicotinate the N-Isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

15                      <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.32 (d, 6H), 2.63 (m, 3H), 4.30 (m, 1H), 7.25 (d, 1H), 7.45-7.51 (m, 2H), 7.63 (s, 1H), 7.69 (t, 1H), 7.77 (d, 1H), 7.82 (dd, 1H), 8.72 (m, 1H), 8.78 (s, 1H), 8.85 (d, 1H), 9.08 (s, 1H), 9.68 (br, NH).

20

**EXAMPLE 29**

**N-Cyclopropyl-1-[3-(1-oxidopyrimidin-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



### Step 1: 5-Bromo-1-oxidopyrimidine

To 5-bromopyrimidine (2.05g, 12.9mmol) in methylene chloride (25ml) was added m-chloroperoxybenzoic acid (ca 70% pure, 3.17g, 12.9mmol) and the resulting mixture was stirred at room temperature for 5 days. Calcium hydroxide (1g) was added and after 10 minutes the mixture was filtered through celite. The product from evaporation of the filtrate was chromatographed on silica gel eluting with ethyl acetate to afford the 5-bromo-1-oxidopyrimidine compound as a white solid.


### Step 2: N-Cyclopropyl-1-[3-(1-oxidopyrimidin-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of **Example 24** but substituting 5-bromo-1-oxidopyrimidine from **Step 1** for ethyl 5-bromonicotinate and N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(1-oxidopyrimidin-5-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.66 (m, 2H), 0.84 (m, 2H), 2.97 (m, 1H), 7.48 (m, 1H), 7.58 (d, 1H), 7.65 (s, 1H), 7.71 (d, 1H), 7.77 (t, 1H), 8.46 (s, 1H), 8.60 (s, 1H), 8.68 (brs, 1H), 8.81 (dd, 1H), 8.98 (s, 1H), 9.02 (s, 1H), 9.72 (br, NH).

## **EXAMPLE 30**

**1-{3-[6-(1-Hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine**

5                   To a suspension of 2,5-dibromopyridine in toluene (12ml/mmol) cooled to  $-78^{\circ}\text{C}$  was added n-butyllithium 2.5M in hexanes (1.05eq) and the resulting mixture was stirred in the cold for 2.5 hours. Acetone (2eq) was added and stirring was continued for 1.5h. After quenching with saturated aqueous ammonium chloride solution, the mixture was warmed to room temperature and partitioned between ethyl acetate and water. The product from the organic phase was chromatographed on silica gel eluting with 20% ethyl acetate in hexane to afford the 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine compound as a syrup.

**Step 2: 5-Bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide**

15                   To a solution of 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine from Step 1 in methylene chloride (5ml/mmol) at room temperature was added m-chloroperoxybenzoic acid 70% (1.1eq) and the resulting mixture was stirred at room temperature for 18 hours. An excess of calcium hydroxide was added and after 5 minutes the mixture was filtered through a bed of celite. The crude product from evaporation of the filtrate was chromatographed on silica gel eluting with 80% ethyl acetate in hexane and the 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide compound was obtained as a white solid.


**Step 3: 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 4** of **Example 1** but substituting 28% aqueous ammonium hydroxide for isopropylamine the 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

**Step 4: 1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Example 24** but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from **Step 2** above for ethyl 5-bromonicotinate and 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the 1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.76 (s, 6H), 5.83 (br, 1H, NH), 7.50 (d, 1H), 7.55 (m, 1H), 7.57-7.62 (m, 2H), 7.65 (m, 2H), 7.72-7.78 (m, 2H), 8.55 (s, 1H, OH), 8.75 (m, 1H), 8.90 (dd, 1H), 9.08 (s, 1H), 9.52 (br, 1H, NH).

**EXAMPLE 31****N-Isopropyl-1-{3-[4-(pyridin-3-yl)phenyl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

**Step 1: N-Isopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

A mixture of N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Step 4** of **Example 1**, diboron pinacol ester (1.1eq), potassium acetate (3.5eq) and [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) in N,N-dimethylformamide (5ml/mmol) was stirred at 85°C for 18 hours. A further amount of diboron pinacol ester (0.4eq) and palladium catalyst (0.05eq) were added and heating and stirring were continued for a further 24 hours. After cooling, the mixture was partitioned between ethyl acetate and water, and the crude product from the organic phase was chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and hexane. The product was then stirred in hexane at room temperature for several hours and filtered to afford the N-Isopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a white solid.

**Step 2: 3-(4-Bromophenyl)pyridine**

A mixture of pyridine-3-boronic acid 1,3-propanediol cyclic ester, 4-bromiodobenzene (1.1eq), [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) and 2M aqueous sodium carbonate (5eq) in N,N-dimethylformamide (2ml/mmol) was stirred at 85°C for 4 hours. After quenching with saturated aqueous ammonium chloride solution, the mixture was partitioned between ethyl acetate and water, and the crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ethyl acetate and hexane to afford the 3-(4-Bromophenyl)pyridine compound as a solid.

**Step 3: N-Isopropyl-1-[3-[4-(pyridin-3-yl)phenyl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


A mixture of the boronate from **Step 1**, 3-(4-bromophenyl)pyridine from **Step 2** (1.5eq), [1,1'-bis (diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq) and 2M aqueous sodium carbonate (5eq) in N,N-dimethylformamide (7ml/mmol) was stirred at 85°C for 1 hour. After cooling, the mixture was

partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 7:3 mixture of ethyl acetate and methylene chloride to afford the N-Isopropyl-1-{3-[4-(pyridin-3-yl)phenyl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.30 (d, 6H), 4.25 (m, 1H), 7.35 (m, 1H), 7.39-7.48 (m, 2H), 7.60-7.75 (m, 6H), 7.80 (d, 1H), 7.90 (d, 1H), 8.58 (d, 1H), 8.70 (m, 1H), 8.82 (d, 1H), 8.88 (s, 1H), 9.08 (s, 1H), 9.68 (br, NH).

### EXAMPLE 32

#### N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



#### Step 1: N-Cyclopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of **Step 1** of **Example 31** but substituting N-cyclopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from step 1 of example 14 for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide the N-Cyclopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

#### Step 2: 3-Bromo-5-methylsulfonylpyridine

To 3,5-dibromopyridine (2.96g, 12.5mmol) in diethyl ether (70ml) at -78°C was added n-butyllithium 1.6M in hexanes (8.6ml, 13.7mmol) and the resulting mixture was stirred in the cold for 3 hours. Dimethyl disulfide (1.12ml, 12.5mmol) was added and the mixture was warmed to room temperature, then partitioned between ether and water. To the crude product from evaporation of the organic phase was added tetrahydrofuran (80ml), methanol (20ml), oxone (17g) and enough saturated aqueous sodium bicarbonate to afford a slightly basic medium. After stirring for 4 hours at room temperature, an excess of 1M aqueous sodium metabisulfite was added, the organic solvents were evaporated, and the residue was partitioned between ethyl acetate and water. The crude product from the organic phase was stirred in a small volume of ethyl acetate and filtered to afford the 3-Bromo-5-methylsulfonylpyridine compound as a solid.

**Step 3: N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


Following the procedure of Step 3 of Example 31 but substituting 3-bromo-5-methylsulfonylpyridine from Step 2 above for 3-(4-bromophenyl)pyridine, and N-cyclopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 1 for N-isopropyl-1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.71 (m, 2H), 0.90 (m, 2H), 3.03 (m, 1H), 3.21 (s, 3H), 7.53 (m, 1H), 7.60 (d, 1H), 7.74 (s, 1H), 7.80 (t, 1H), 7.86 (d, 1H), 8.45 (m, 1H), 8.74 (m, 1H), 8.86 (d, 1H), 9.09 (s, 1H), 9.20 (d, 2H), 9.78 (br, NH).

**EXAMPLE 33**

**N-Cyclopropyl-1-[3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





**Step 1: Methyl 2-bromoisonicotinate**

To a solution of 2-bromoisonicotinic acid (*Chem. Pharm. Bull.*, 38:2446(1990)) (2.0g) in tetrahydrofuran (100ml) was added excess ethereal  
 5 diazomethane and the resulting mixture was stirred at room temperature for 1 hour. The mixture was evaporated and the product chromatographed on silica gel eluting with a 1:3 mixture of ethyl acetate and hexane to afford the Methyl 2-bromoisonicotinate ester as a colorless liquid.

10 **Step 2: 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine**

Following the procedure of **Step 1** of **Example 25**, but substituting methyl 2-bromoisonicotinate from **Step 1** for ethyl 5-bromonicotinate, the 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine compound was obtained as a white solid.

15 **Step 3: 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide**

Following the procedure of **Step 2** of **Example 30** but substituting 2-bromo-4-(1-hydroxy-1-methylethyl)pyridine from **Step 2** for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine the 2-Bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide compound was obtained as a white solid.

20


**Step 4: N-Cyclopropyl-1-{3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3** of **Example 32**, but substituting 2-bromo-4-(1-hydroxy-1-methylethyl)pyridine-N-oxide from **Step 3** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[4-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.57 (m, 2H), 0.79 (m, 2H), 1.45 (s, 6H), 2.90 (m, 1H), 5.35 (s, 1H, OH), 7.48 (m, 1H), 7.64 (m, 1H), 7.72 (m, 3H), 8.11 (m, 2H), 8.30 (d, 1H), 8.72 (dd, 1H), 8.78 (m, 1H), 8.82 (s, 1H), 9.72 (br, NH).

#### EXAMPLE 34

**N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl} 1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



##### Step 1: 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine

A solution of 2,5-dibromopyridine in diethyl ether (5ml/mmol) was cooled to  $-78^\circ\text{C}$ , and n-butyllithium 2.5M in hexanes (1.05eq) was added slowly. After 2h in the cold, acetone (1.3eq) was added and stirring was continued for 1 hour. The resulting mixture was quenched with saturated aqueous ammonium chloride solution, warmed to room temperature, and partitioned between ether and water. The crude product from the organic phase was triturated with 1:1 ether-hexane and filtered to afford the 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine compound as a solid.


**Step 2:** N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of **Step 3** of **Example 32**, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.71 (m, 2H), 0.90 (m, 2H), 1.68 (s, 6H), 1.85 (s, 1H, OH), 3.04 (m, 1H), 7.45-7.52 (m, 2H), 7.71 (t, 1H), 7.79 (d, 1H), 7.95 (dd, 1H), 8.16 (s, 1H), 8.20 (d, 1H), 8.72 (m, 1H), 8.80-8.87 (m, 2H), 9.12 (s, 1H), 9.82 (br, NH).

**EXAMPLE 35**

**N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1:** 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine

Following the sequence described in **Steps 1-2** of **Example 33**, but substituting 4-bromopicolinic acid (*Aust. J. Chem.* 24:390(1971)) for 2-bromoisonicotinic acid in **Step 1**, the 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine compound was obtained as a white solid.

**Step 2: N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


Following the procedure of **Step 3** of **Example 32**, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine from **Step 1** for 3-bromo-5-

- 5 methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)pyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.78 (m, 2H), 1.48 (s, 6H), 2.91 (m, 1H), 5.27 (s, 1H, OH), 7.62-7.66 (m, 2H), 7.72-7.79 (m, 2H), 8.01 (m, 1H), 8.10 (s, 1H), 8.58 (d, 1H), 8.73-8.79 (m, 2H), 8.84 (s, 1H), 9.73 (br, NH).

**EXAMPLE 36**

- 15 **Synthesis of N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine N-oxide**

- Following the procedure of **Step 2** of **Example 30**, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine from **Step 1** of **Example 35** for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the 4-Bromo-2-(1-hydroxy-1-methylethyl)pyridine N-oxide compound was obtained as a white solid.


- 25 **Step 2: N-Cyclopropyl-1-{3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3 of Example 32**, but substituting 4-bromo-2-(1-hydroxy-1-methylethyl)pyridine-N-oxide from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-[3-(1-hydroxy-1-methylethyl)-1-oxidopyridin-4-yl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a beige solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.57 (m, 2H), 0.78 (m, 2H), 1.62 (s, 6H), 2.90 (m, 1H), 6.99 (s, 1H, OH), 7.65-7.84 (m, 4H), 7.94 (s, 1H), 8.03 (dd, 1H), 8.15 (s, 1H), 8.38 (d, 1H), 8.73-8.78 (m, 2H), 8.83 (s, 1H), 9.73 (br, NH).

### EXAMPLE 37

#### N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



#### Step 1: 5-Bromo-2-isopropylthiopyridine

To a mixture of 2,5-dibromopyridine (2.07g, 8.73mmol) and 2-propanethiol (0.97ml, 10.4mmol) in N,N-dimethylformamide (20ml) at 0°C was added portionwise sodium hydride 60% dispersed in oil (450mg, 11.3mmol). The resulting mixture was stirred at room temperature for 1 hour, then partitioned between ether and water. The crude product from the organic phase was chromatographed on silica gel eluting with 10% ethyl acetate in hexane to afford the 5-Bromo-2-isopropylthiopyridine compound as a solid.

**Step 2: 5-Bromo-2-isopropylsulfonylpyridine**

To a solution of 5-bromo-2-isopropylthiopyridine from **Step 1** (2.03g, 8.75mmol) in tetrahydrofuran (50ml) and methanol (25ml) at 0°C was added oxone (15.8g, 25.8mmol) and then saturated aqueous sodium bicarbonate (25ml). The resulting mixture was stirred at room temperature for 6 hours. The mixture was quenched with aqueous sodium bicarbonate and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with 20% ethyl acetate in hexane to afford the 5-Bromo-2-isopropylsulfonylpyridine compound as a white solid.


**Step 3: N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3** of **Example 32**, but substituting 5-bromo-2-isopropylsulfonylpyridine from **Step 2** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-isopropylsulfonylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.70 (m, 2H), 0.89 (m, 2H), 1.39 (d, 6H), 3.00 (m, 1H), 3.82 (m, 1H), 7.51 (m, 1H), 7.60 (d, 1H), 7.72 (s, 1H), 7.80 (t, 1H), 7.83 (d, 1H), 8.15-8.24 (m, 2H), 8.72 (m, 1H), 8.86 (dd, 1H), 9.03 (s, 1H), 9.10 (s, 1H), 9.77 (br, NH).

25

**EXAMPLE 38****N-Cyclopropyl-1-[3-(6-methoxypyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

**Step 1: 5-Bromo-2-methoxypyridine**


To a solution of 2,5-dibromopyridine (6.95g, 29mmol) in N,N-dimethylformamide (5ml) was added methanol (3.56ml) and 1M potassium tert-butoxide (32.3ml) and the resulting mixture was stirred at room temperature for 18 hours. The resulting slurry was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ether and hexane to afford the 5-Bromo-2-methoxypyridine compound as an oil.

**Step 2: N-Cyclopropyl-1-[3-(6-methoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of Step 3 of Example 32, but substituting 5-bromo-2-methoxypyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-cyclopropyl-1-[3-(6-methoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.71 (m, 2H), 0.89 (m, 2H), 3.00 (m, 1H), 4.00 (s, 3H), 6.85 (d, 1H), 7.44 (d, 1H), 7.50 (m, 1H), 7.62 (s, 1H), 7.68 (t, 1H), 7.73 (d, 1H), 7.83 (dd, 1H), 8.44 (s, 1H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.82 (br, NH).

**EXAMPLE 39****N-Cyclopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 3 of Example 32**, but substituting 5-bromo-2-methylpyridine from **Step 1 of Example 28** for 3-bromo-5-methylsulfonylpyridine, the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 2.65 (s, 3H), 3.03 (m, 1H), 7.28 (d, 1H), 7.45-7.53 (m, 2H), 7.66 (s, 1H), 7.72 (t, 1H), 7.80 (d, 1H), 7.84 (dd, 1H), 8.73 (m, 1H), 8.80 (s, 1H), 8.86 (dd, 1H), 9.11 (s, 1H), 9.82 (br, NH).

10

**EXAMPLE 40**

**N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





**Step 1: 5-Bromo-2-(2,2,2-trifluoroethoxy)pyridine**

Following the procedure of **Step 1** of **Example 38**, but substituting 2,2,2-trifluoroethanol for methanol, with heating at 70°C for 18 hours, the 5-Bromo-2-(2,2,2-trifluoroethoxy)pyridine compound was obtained as an oil.

5

**Step 2: N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


Following the procedure of **Step 3** of **Example 32**, but substituting 5-bromo-2-(2,2,2-trifluoroethoxy)pyridine from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[6-(2,2,2-trifluoroethoxy)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 3.03 (m, 1H), 4.85 (q, 2H), 7.00 (d, 1H), 7.43-7.53 (m, 2H), 7.62 (s, 1H), 7.69-7.78 (m, 2H), 7.92 (dd, 1H), 8.42 (s, 1H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.80 (br, NH).

15

**EXAMPLE 41**

20

**N-Cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3** of **Example 32**, but substituting 3,5-dibromopyridine for 3-bromo-5-methylsulfonylpyridine, the title compound was obtained as a white solid.


25

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.58 (m, 2H), 0.79 (m, 2H), 2.90 (m, 1H), 7.65 (m, 1H), 7.71-7.77 (m, 2H), 8.03 (d, 1H), 8.14 (s, 1H), 8.49 (s, 1H), 8.74 (brs, 1H), 8.79 (brs, 1H), 8.86 (s, 1H), 9.01 (s, 1H), 9.73 (br, NH).

5

### EXAMPLE 42

#### N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



10

#### Step 1: 2-Benzyloxy-5-bromopyridine

A mixture of 2,5-dibromopyridine, benzyl alcohol (1.3eq), potassium hydroxide pellets (2.4eq) and dibenzo-18-crown-6 (0.05eq) in toluene (4ml/mmol) was refluxed with azeotropic removal of water for 3 hours. After evaporation of the toluene, the resulting mixture was partitioned between chloroform and water. The crude product from the organic phase was recrystallized from ether-hexane to afford the 2-Benzyloxy-5-bromopyridine compound as a solid.

15

#### Step 2: N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

20


Following the procedure of Step 3 of Example 32, but substituting 2-benzyloxy-5-bromopyridine from Step 1 for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-benzyloxy-3-pyridin-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.79 (m, 2H), 2.91 (m, 1H), 5.42 (s, 2H) 7.00 (d, 1H), 7.32-7.48 (m, 5H), 7.61-7.72 (m, 3H), 7.90 (d, 1H), 7.99 (s, 1H), 8.14 (d, 1H), 8.59 (s, 1H), 8.73-8.84 (m, 3H), 9.73 (br, NH).

5

### EXAMPLE 43

**N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



10

**Step 1: 5-Bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide**

Following the procedure of **Steps 1 and 2 of Example 30**, but substituting dicyclopropyl ketone for acetone in **Step 1**, the 5-Bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide compound was obtained as a solid.

15

**Step 2: N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3 of Example 32**, but substituting 5-bromo-2-dicyclopropyl(hydroxy)methylpyridine N-oxide from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[6-dicyclopropyl(hydroxy)methyl-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a white solid.

20


<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.52 (m, 4H), 0.70 (m, 4H), 0.76 (m, 2H), 0.89 (m, 2H), 1.35 (m, 2H), 3.02 (m, 1H), 7.52 (m, 1H), 7.58 (m, 1H), 7.62 (dd, 1H), 7.68

(s, 1H), 7.73-7.80 (m, 3H), 8.15 (br, 1H, OH), 8.49 (s, 1H), 8.72 (m, 1H), 8.85 (dd, 1H), 9.09 (s, 1H), 9.78 (br, NH).

5

#### EXAMPLE 44

**N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



#### 10 Step 1: 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide

Following the procedure of **Step 2 of Example 30**, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine from **Step 1 of Example 34** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the 2-Bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide compound was obtained as a white solid.

15

#### Step 2: N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

Following the procedure of **Step 3 of Example 32**, but substituting 2-bromo-5-(1-hydroxy-1-methylethyl)pyridine N-oxide from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.


20

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.69 (m, 2H), 0.88 (m, 2H), 1.63 (s, 6H), 2.20 (s, 1H, OH), 2.98 (m, 1H), 7.38-7.49 (m, 3H), 7.52 (d, 1H), 7.70 (t, 1H), 7.98-8.04 (m, 2H), 8.50 (s, 1H), 8.69 (m, 1H), 8.80 (dd, 1H), 9.08 (s, 1H), 9.75 (br, NH).

5

**EXAMPLE 45**

**N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



10

Following the procedure of **Step 3** of **Example 32**, but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine from **Step 1** of **Example 30** for 3-bromo-5-methylsulfonylpyridine, the title compound was obtained as a solid.


<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.62 (s, 6H), 3.02 (m, 1H), 4.85 (s, 1H, OH), 7.48-7.53 (m, 3H), 7.68 (s, 1H), 7.73 (t, 1H), 7.80 (d, 1H), 7.95 (dd, 1H), 8.72 (m, 1H), 8.81 (s, 1H), 8.86 (dd, 1H), 9.10 (s, 1H), 9.78 (br, NH).

15

**EXAMPLE 46**

20

**N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 5-(3-Aminophenyl)- 2-(1-hydroxy-1-methylethyl) pyridine**

Following the procedure of **Step 5** of **Example 1**, but substituting 3-aminophenylboronic acid for 3-acetyl phenylboronic acid and 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine from **Step 1** of **Example 30** for N-isopropyl-1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the 5-(3-Aminophenyl)- 2-(1-hydroxy-1-methylethyl) pyridine compound was obtained as a solid.

**Step 2: 1-{3-[6-(1-Hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid**

Following the procedures of **Steps 1-3** of **Example 1**, but substituting 5-(3-aminophenyl)- 2-(1-hydroxy-1-methylethyl)pyridine for 3-bromoaniline from **Step 1** in the **First Step**, the 1-{3-[6-(1-Hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a solid.

**Step 3: N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 4** of **Example 1**, but substituting the acid from **Step 2** for 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid, and isobutylamine for isopropylamine, the N-Isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a cream-colored solid.


<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.98 (d, 6H), 1.53 (s, 6H), 1.88 (m, 1H), 3.26 (t, 2H), 4.66 (s, 1H, OH), 7.60 (m, 1H), 7.69 (d, 1H), 7.76-7.79 (m, 2H), 7.95 (d, 1H), 8.05 (s, 1H), 8.16 (dd, 1H), 8.73 (m, 1H), 8.79 (dd, 1H), 8.90 (s, 1H), 8.94 (s, 1H), 9.83 (br, NH).

5

### EXAMPLE 47

N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-  
1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide

10



**Step 1: 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid**

Following the procedures of Steps 1-3 of Example 1, but substituting 3,5-dibromoaniline for 3-bromoaniline in Step 1, the 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid compound was obtained as a beige solid.

**Step 2: N-Cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

20

Following the procedure of Step 4 of Example 1, but substituting the 1-(3,5-Dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid from Step 1 for 1-(3-bromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxylic acid, and cyclopropylamine for isopropylamine, the N-Cyclopropyl-1-(3,5-

dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

**Step 3: 2-(1-Hydroxy-1-methylethyl)-5-tributylstannylpyridine**

- 5 To a suspension of 2,5-dibromopyridine in toluene (5ml/mmol) at -78°C, was added n-butyllithium 2.5M in hexanes (1eq) and the resulting mixture was stirred in the cold for 2.5 hours. Acetone (1eq) was added, and the mixture was warmed to -50°C and became a brown solution. After cooling down to -78°C, more n-butyllithium (1eq) was added along with ether (2ml/mmol). After stirring in the cold for a further hour, tributyltin chloride (1.1eq) was added and the mixture was warmed to room temperature and stirred for 2 hours. The mixture was quenched with saturated aqueous ammonium chloride solution and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:9 mixture of ethyl acetate and hexane to afford the 2-(1-Hydroxy-1-methylethyl)-5-tributylstannylpyridine compound as a colorless liquid.
- 10
- 15

**Step 4: N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


- A mixture of N-cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from Step 2, 2-(1-hydroxy-1-methylethyl)-5-tributylstannylpyridine from Step 3 (1.4eq), 1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.05eq), and cuprous iodide (0.05eq) in N,N-dimethylformamide (15ml/mmol) was stirred at 85°C for 5 hours. After cooling the resulting mixture was partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 1:6:3 mixture of ethanol, ethyl acetate and methylene chloride to afford the N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound as a solid.
- 20
- 25

- <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.62 (s, 6H), 3.02 (m, 1H), 4.76 (s, 1H, OH), 7.50-7.56 (m, 2H), 7.62 (s, 1H), 7.69 (s, 1H), 7.90-7.96 (m, 2H), 8.74 (m, 1H), 8.79 (s, 1H), 8.86 (dd, 1H), 9.07 (s, 1H), 9.74 (br, NH).
- 30

**EXAMPLE 48**



**N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**5    Step 1: 2-(1-Hydroxy-1-methylethyl)-6-tributylstannylpyridine**

Following the procedure of **Step 3** of **Example 47**, but substituting 2,6-dibromopyridine for 2,5-dibromopyridine, the 2-(1-Hydroxy-1-methylethyl)-6-tributylstannylpyridine compound was obtained.


**10    Step 2: N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 4** of **Example 47**, but substituting 2-(1-hydroxy-1-methylethyl)-6-tributylstannyl pyridine from **Step 1** for 2-(1-hydroxy-1-methylethyl)-5-tributylstannylpyridine, the N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-2-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

**15    <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.61 (s, 6H), 3.04 (m, 1H), 5.13 (s, 1H, OH), 7.40 (d, 1H), 7.46-7.53 (m, 2H), 7.70-7.76 (m, 2H), 7.85 (t, 1H), 8.13 (s, 1H), 8.22 (d, 1H), 8.73 (m, 1H), 8.87 (d, 1H), 9.12 (s, 1H), 9.83 (br, 20    NH).**

**EXAMPLE 49**

**N-Isopropyl-1-[3-(4-methylsulfonylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




To a mixture of N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 22** in tetrahydrofuran (24ml/mmol), methanol (12ml/mmol), and water (12ml/mmol), was added oxone (2.24eq) and the resulting mixture was stirred at room temperature for 2 hours. The mixture was quenched with saturated aqueous sodium bicarbonate and partitioned between ethyl acetate and water. The crude product from the organic phase was chromatographed on silica gel eluting with 30% ether in methylene chloride to afford the title compound as a white solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 1.25 (d, 6H), 3.16 (s, 3H), 4.18 (m, 1H), 7.60 (m, 1H), 7.74 (d, 1H), 7.79 (t, 1H), 7.99 (d, 1H), 8.05 (s, 4H), 8.09 (s, 1H), 8.72 (m, 1H), 8.78 (dd, 1H), 8.93 (s, 1H), 9.64 (br, NH).

**EXAMPLE 50**

**N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



**Step 1: 5-Bromo-2-methylthiopyridine**

A mixture of 2,5-dibromopyridine and sodium thiomethoxide (1.3eq) in N,N-dimethylformamide (2ml/mmol) was stirred at room temperature for 20 minutes then cooled to 0°C. After diluting with cold water the precipitate was filtered to afford the 5-Bromo-2-methylthiopyridine compound as a solid.

**Step 2: N-Cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Step 3** of **Example 32**, but substituting 5-bromo-2-methylthiopyridine from **Step 1** for 3-bromo-5-methylsulfonylpyridine, the N-Cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.


**Step 3: N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

Following the procedure of **Example 49**, but substituting N-cyclopropyl-1-[3-(6-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Step 2** for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the N-Cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.66 (m, 2H), 0.84 (m, 2H), 2.97 (m, 1H), 3.26 (s, 3H), 7.48 (m, 1H), 7.55 (d, 1H), 7.67 (s, 1H), 7.74-7.80 (m, 2H), 8.14-8.19 (m, 2H), 8.68 (m, 1H), 8.81 (dd, 1H), 8.96 (s, 1H), 9.05 (s, 1H), 9.73 (br, NH).

## EXAMPLE 51

5                    **N-Isopropyl-1-[3-(5-methylsulfonylpyridin-3-yl)phenyl]-1,4-**  
                     **dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Example 49**, but substituting N-isopropyl-  
1-[3-(5-methylthiopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-  
10 carboxamide from **Example 15** for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-  
1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained  
as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.33 (d, 6H), 3.20 (s, 3H), 4.31 (m, 1H), 7.52 (m,  
1H), 7.60 (d, 1H), 7.73 (s, 1H), 7.79 (t, 1H), 7.86 (d, 1H), 8.48 (m, 1H), 8.73 (m, 1H),  
15 8.88 (d, 1H), 9.08 (s, 1H), 9.19 (d, 2H), 9.68 (br, NH).

## EXAMPLE 52

20                    **N-Cyclopropyl-1-[3-(4-ethylsulfonylphenyl)phenyl]-1,4-**  
                     **dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Example 49**, but substituting N-cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 18** for N-isopropyl-1-[3-(4-methylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.90 (m, 2H), 1.35 (t, 3H), 3.02 (m, 1H), 3.18 (q, 2H), 7.48-7.56 (m, 2H), 7.70 (s, 1H), 7.75 (t, 1H), 7.84 (m, 3H), 8.03 (d, 2H), 8.73 (m, 1H), 8.85 (dd, 1H), 9.10 (s, 1H), 9.80 (br, NH).

### EXAMPLE 53

**N-Cyclopropyl-1-[3-(4-ethylsulfinylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




To a solution of N-cyclopropyl-1-[3-(4-ethylthiophenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 18**, in a 1:1 mixture of methylene chloride and methanol (9ml/mmol), was added at 0°C magnesium  
 5 monoperoxyphthalate hexahydrate (MMPP, 0.5molareq) and the resulting mixture was stirred in the cold for 2 hours. The mixture was quenched with saturated aqueous sodium bicarbonate and partitioned between methylene chloride and water. The crude product from the organic phase was chromatographed on silica gel eluting with a 90:9:1 mixture of methylene chloride, ethanol and 28% aqueous ammonium  
 10 hydroxide to afford the title compound as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.68 (m, 2H), 0.85 (m, 2H), 1.15 (m, 3H), 2.80 (m, 1H), 2.94 (m, 1H), 2.98 (m, 1H), 7.45-7.50 (m, 2H), 7.65-7.73 (m, 4H), 7.76-7.82 (m, 3H), 8.71 (m, 1H), 8.83 (dd, 1H), 9.06 (s, 1H), 9.78 (br, NH).

15

#### EXAMPLE 54

**N-Isopropyl-1-{3-[4-(1-oximidoethyl)phenyl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




To a solution of N-isopropyl-1-[3-(4-acetylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 4** in pyridine (11ml/mmol) at room temperature was added hydroxylamine hydrochloride (2.1eq) and the resulting mixture was stirred for 16 hours. The mixture was filtered through celite and the filtrate evaporated. The residue was dissolved in ethyl acetate, washed with saturated aqueous sodium carbonate and then water, dried and evaporated. The residue was stirred in a small volume of acetone and filtered to afford the title compound as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 2.27 (s, 3H), 4.30 (m, 1H), 7.39 (d, 1H), 7.46 (m, 1H), 7.56 (d, 2H), 7.59-7.63 (m, 2H), 7.66 (d, 2H), 7.72 (d, 1H), 8.17 (s, 1H, OH), 8.69 (brs, 1H), 8.82 (d, 1H), 9.10 (s, 1H), 9.71 (br, NH).

#### EXAMPLE 55

**N-Isopropyl-1-{3-[ 4-(4-piperazin-1-yl)phenyl]-phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




To a solution of N-isopropyl-1-[3-[4-(4-tertbutyloxycarbonylpiperazin-1-yl)phenyl]-phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 11** in methylene chloride (10ml/mmol) was added trifluoroacetic acid (6ml/mmol) and the resulting mixture was stirred at room temperature for 2 hours, then warmed gently for 15 minutes. The mixture was evaporated and the crude product was chromatographed on silica gel eluting with a 9:0.9:0.1 mixture of methylene chloride, methanol and 28% aqueous ammonium hydroxide to afford the title compound as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 2.99 (m, 4H), 3.16 (m, 4H), 4.25 (m, 1H), 6.94 (d, 2H), 7.29 (d, 1H), 7.42 (m, 1H), 7.50 (d, 2H), 7.52-7.58 (m, 2H), 7.69 (d, 1H), 8.66 (m, 1H), 8.78 (dd, 1H), 9.04 (s, 1H), 9.69 (br, NH).

## EXAMPLE 56

**N-Cyclopropyl-1-[3-(4-methylsulfonylmethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**





**Step 1: N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**


A mixture of N-cyclopropyl-1-[3-(4-hydroxymethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 16**, carbon tetrabromide (2eq), and diphos (0.6 molareq) in methylene chloride (15ml/mmol) was stirred at room temperature for 3 hours. The mixture was concentrated at room temperature and chromatographed on silica gel eluting with a 1:1 mixture of ethyl acetate and methylene chloride to afford the N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide compound.

**Step 2: N-Cyclopropyl-1-[3-(4-methylsulfonylmethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

To a solution of N-Cyclopropyl-1-[3-(4-bromomethylphenyl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Step 1** in N,N-dimethylformamide (20ml/mmol) was added methanesulfinic acid sodium salt (1.3eq) and the resulting mixture was stirred at room temperature for 18 hours. To the mixture was added saturated aqueous ammonium chloride solution and ethyl acetate, and the insoluble solid was filtered and washed well with water, hexane, ether and ethyl acetate to afford the title compound as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.72 (m, 2H), 0.89 (m, 2H), 2.85 (s, 3H), 3.04 (m, 1H), 4.34 (s, 2H) 7.46-7.52 (m, 2H), 7.55 (d, 2H), 7.65-7.73 (m, 4H), 7.80 (d, 1H), 8.76 (m, 1H), 8.85 (d, 1H), 9.12 (s, 1H), 9.82 (br, NH).

## EXAMPLE 57

**N-Cyclopropyl-1-[3-(1,6-dihydro-6-oxopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

5

To a solution of N-cyclopropyl-1-[3-(6-benzyloxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 42** in 1,2-dichloroethane (25ml/mmol) was added trifluoroacetic acid (1.5ml/mmol) and the resulting mixture was stirred at 60°C for 18 hours. More trifluoroacetic acid was added (0.75ml/mmol) and heating was continued for a further 24 hours. The cooled mixture was diluted with methylene chloride and saturated aqueous sodium bicarbonate was added, resulting in precipitation of a solid which was filtered. From the filtrate the organic phase was collected and evaporated to a solid which was combined with the previous filtered solid. This mixture was chromatographed on silica gel eluting with 10% methanol in methylene chloride to afford the title compound as a white fluffy solid.


15

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.90 (m, 1H), 6.45 (d, 1H), 7.52 (m, 1H), 7.61-7.65 (m, 2H), 7.78 (d, 1H), 7.85 (s, 1H), 7.89-7.93 (m, 2H), 8.74 (d, 1H), 8.78-8.81 (m, 2H), 9.73 (br, NH), other NH >11ppm.

20

## EXAMPLE 58

**N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]pyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 4** of **Example 47**, but substituting N-cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 41** for N-cyclopropyl-1-(3,5-dibromophenyl)-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a white solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.58 (m, 2H), 0.79 (m, 2H), 2.91 (m, 1H), 5.30 (s, 1H, OH), 7.65 (m, 1H), 7.71-7.79 (m, 3H), 8.12 (d, 1H), 8.23-8.26 (m, 2H), 8.49 (s, 1H), 8.75 (dd, 1H), 8.80 (m, 1H), 8.87 (s, 1H), 8.97 (m, 2H), 9.04 (s, 1H), 9.74 (br, NH).

15

**EXAMPLE 59**

**N-Isopropyl-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2 of Example 30**, but substituting N-isopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 7** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  1.21 (d, 6H), 4.10 (m, 1H), 7.51 (t, 1H), 7.64 (m, 1H), 7.71-7.75 (m, 3H), 7.97 (m, 1H), 8.09 (s, 1H), 8.23 (d, 1H), 8.69-8.77 (m, 3H), 8.84 (s, 1H), 9.66 (br, NH).

#### EXAMPLE 60

**N-(2,6-Dichloropyridin-4-yl)-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2 of Example 30**, but substituting N-(2,6-dichloropyridin-4-yl)-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-

one-3-carboxamide from **Example 10** for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the title compound was obtained as a white solid.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  7.51 (m, 1H), 7.69-7.78 (m, 4H), 7.99 (dd, 1H), 8.14 (s, 1H), 8.24 (dd, 1H), 8.70 (s, 1H), 8.73 (s, 2H), 8.84 (m, 2H), 8.99 (s, 1H), 12.05 (br, NH).

### EXAMPLE 61

**N-Isopropyl-1-[3-(5-carboethoxy-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2** of **Example 30**, but substituting N-isopropyl-1-[3-(5-carboethoxypyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 24** for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the title compound was obtained as a white solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.28 (d, 6H), 1.40 (t, 3H), 4.28 (m, 1H), 4.43 (q, 2H), 7.49 (dd, 1H), 7.56 (m, 1H), 7.68 (s, 1H), 7.73 (d, 2H), 8.04 (s, 1H), 8.60 (s, 1H), 8.68 (dd, 1H), 8.77 (s, 1H), 8.82 (d, 1H), 9.01 (s, 1H), 9.61 (br, NH).

### EXAMPLE 62

**N-Isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2 of Example 30**, but substituting N-isopropyl-1-{3-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 25** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.29 (d, 6H), 1.60 (s, 6H), 4.11 (brs, 1H), 4.23 (m, 1H), 7.42-7.51 (m, 2H), 7.58 (s, 2H) 7.65 (m, 2H), 8.28 (s, 1H), 8.33 (s, 1H), 8.64 (m, 1H), 8.80 (d, 1H), 8.98 (s, 1H), 9.61 (br, NH).

**EXAMPLE 63**

**N-Isopropyl-1-{3-[6-(2-methylpropyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



Following the procedure of **Step 2** of **Example 30**, but substituting N-isopropyl-1-[3-[6-(2-methylpropyl)pyridin-3-yl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 26**

- 5 for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as an off- white solid.


$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.98 (d, 6H), 1.29 (d, 6H), 2.29 (m, 1H), 2.32 (d, 2H), 4.26 (m, 1H), 7.28 (d, 1H), 7.38 (d, 1H), 7.47-7.52 (m, 2H), 7.60 (s, 1H), 7.69 (m, 2H), 8.53 (s, 1H), 8.69 (m, 1H), 8.82 (dd, 1H), 9.03 (s, 1H), 9.62 (br, NH).

10

#### EXAMPLE 64

**N-Isopropyl-1-[3-(6-methyl-1-oxopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

15




Following the procedure of **Step 2** of **Example 30**, but substituting N-isopropyl-1-[3-(6-methylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 28** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as an off-white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.32 (d, 6H), 2.60 (s, 3H), 4.30 (m, 1H), 7.35-7.45 (m, 2H), 7.50 (m, 2H), 7.62 (s, 1H), 7.72 (d, 2H), 8.58 (s, 1H), 8.72 (m, 1H), 8.85 (dd, 1H), 9.06 (s, 1H), 9.66 (br, NH).

### EXAMPLE 65

**N-Cyclopropyl-1-[3-(1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**






Following the procedure of **Step 2** of **Example 30**, but substituting N-cyclopropyl-1-[3-(pyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 14** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.78 (m, 2H), 2.90 (m, 1H), 7.52 (t, 1H), 7.65 (m, 1H), 7.72-7.76 (m, 3H), 7.98 (m, 1H), 8.10 (s, 1H), 8.25 (d, 1H), 8.70-8.79 (m, 3H), 8.85 (s, 1H), 9.72 (br, NH).

10

**EXAMPLE 66**

**N-Cyclopropyl-1-{3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



15


Following the procedure of **Step 2** of **Example 29** but substituting 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from **Step 2** of **Example 30** for 5-bromo-1-oxidopyrimidine, the title compound was obtained as a white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.66 (m, 2H), 0.85 (m, 2H), 1.70 (s, 6H), 2.97 (m, 1H), 7.43-7.49 (m, 2H), 7.52-7.56 (m, 2H), 7.61 (s, 2H), 7.71-7.74 (m, 2H), 8.49 (s, 1H), 8.68 (m, 1H), 8.80 (d, 1H), 9.02 (s, 1H), 9.74 (br, NH).

20

**EXAMPLE 67**

**N-Cyclopropyl-1-[3-(1-oxidopyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2 of Example 30**, but substituting N-cyclopropyl-1-[3-(pyridin-4-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 17** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a white solid.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.57 (m, 2H), 0.79 (m, 2H), 2.92 (m, 1H), 7.62-7.70 (m, 2H), 7.75 (t, 1H), 7.88 (d, 2H), 8.03 (d, 1H), 8.15 (s, 1H), 8.30 (d, 2H), 8.75 (d, 1H), 8.80 (m, 1H), 8.86 (s, 1H), 9.73 (br, NH).

**EXAMPLE 68**

**N-Cyclopropyl-1-[3-(5-bromo-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




Following the procedure of **Step 2** of **Example 30**, but substituting N-cyclopropyl-1-[3-(5-bromopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 41** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a light yellow solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.56 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 7.65 (m, 1H), 7.71-7.74 (m, 2H), 8.02-8.06 (m, 2H), 8.15 (s, 1H), 8.60 (s, 1H), 8.73-8.79 (m, 3H), 8.86 (s, 1H), 9.73 (br, NH).

### EXAMPLE 69

**N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]pyridin-3-yl}phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



Following the procedure of **Step 2 of Example 30**, but substituting N-cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]pyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 58** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, and using 1.6eq. of m-chloroperoxybenzoic acid, the title compound was obtained as a white solid.


$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.57 (m, 2H), 0.78 (m, 2H), 2.91 (m, 1H), 6.94 (s, 1H, OH), 7.65 (m, 1H), 7.71-7.79 (m, 3H), 7.97 (dd, 1H), 8.13 (d, 1H), 8.25 (s, 1H), 8.55 (s, 1H), 8.74 (dd, 1H), 8.80 (m, 1H), 8.87 (s, 1H), 8.91 (s, 1H), 9.00 (s, 1H), 9.09 (s, 1H), 9.73 (br, NH).

10

### EXAMPLE 70

**N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-oxidopyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**

15




From the procedure of **Example 69**, the title compound was also obtained as a white solid.

$^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  0.57 (m, 2H), 0.79 (m, 2H), 2.92 (m, 1H), 5.32 (s, 1H, OH), 7.65 (m, 1H), 7.72-7.80 (m, 3H), 8.08-8.17 (m, 2H), 8.27 (m, 2H), 8.70-8.82 (m, 4H), 8.88 (s, 1H), 8.98 (s, 1H), 9.73 (br, NH).

20

### EXAMPLE 71

**N-Cyclopropyl-1-[[3-{5-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]-1-oxidopyridin-3-yl}phenyl]]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




From the procedure of **Example 69** the title compound was also obtained as a white solid.

$^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.58 (m, 2H), 0.80 (m, 2H), 2.92 (m, 1H), 6.85 (brs, 1H, OH), 7.65 (m, 1H), 7.70-7.80 (m, 3H), 7.96 (d, 1H), 8.13 (m, 2H), 8.29 (s, 1H), 8.71-8.84 (m, 4H), 8.89 (s, 1H), 8.92 (s, 1H), 9.73 (br, NH).

#### EXAMPLE 72

**N-Isopropyl-1-[3-(1-oxidoquinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide**




Following the procedure of **Step 2 of Example 30**, but substituting N-isopropyl-1-[3-(quinolin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-carboxamide from **Example 12** for 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine, the title compound was obtained as a solid.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.30 (d, 6H), 4.28 (m, 1H), 7.49 (dd, 1H), 7.54 (d, 1H), 7.66-7.85 (m, 5H), 7.92 (m, 2H), 8.69-8.75 (m, 2H), 8.84 (d, 1H), 8.86 (s, 1H), 9.08 (s, 1H), 9.64 (br, NH).

10

### EXAMPLE 73

**N-Isobutyl-1-[3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**




15

To a mixture of N-isobutyl-1-{3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 46** in 13:1 methylene chloride/methanol (33ml/mmol) at room temperature was added magnesium monoperoxyphthalate hexahydrate (MMPP, 1.1 molar eq) and the resulting mixture was stirred at room temperature for 24 hours. The mixture was filtered through a bed of celite and the filtrate was washed with aqueous sodium carbonate, then water and dried. The crude product was chromatographed on silica gel eluting with 8% ethanol in ethyl acetate and the solid obtained was stirred at room temperature in ether for several hours and filtered to afford the title compound as a light pink solid.

<sup>1</sup>H NMR (Acetone-d<sub>6</sub>) δ 0.98 (d, 6H), 1.61 (s, 6H), 1.88 (m, 1H), 3.26 (t, 2H), 7.52 (s, 1H, OH), 7.61 (m, 1H), 7.66 (d, 1H), 7.77-7.82 (m, 2H), 7.88 (d, 1H), 7.99 (d, 1H), 8.12 (s, 1H), 8.68 (s, 1H), 8.73 (m, 1H), 8.80 (dd, 1H), 8.93 (s, 1H), 9.81 (br, NH).

#### EXAMPLE 74

N-Cyclopropyl-1-[3-(6-methyl-1-oxidopyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide




Following the procedure of **Example 73**, but substituting N-cyclopropyl-1-[3-(6-methylpyridin-3-yl)]phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 39** for N-isobutyl-1-{3-[6-(1-hydroxy-1-

methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide, the title compound was obtained as a solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.72 (m, 2H), 0.91 (m, 2H), 2.61 (s, 3H), 3.02 (m, 1H), 7.38 (d, 1H), 7.45 (dd, 1H), 7.49-7.58 (m, 2H), 7.66 (s, 1H), 7.75 (m, 2H), 8.61 (s, 1H), 8.72 (m, 1H), 8.87 (dd, 1H), 9.08 (s, 1H), 9.78 (br, NH).

#### EXAMPLE 75

##### 10 N-Cyclopropyl-1-[3-(6-methylsulfonyl-1-oxidopyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide



To a suspension of N-cyclopropyl-1-[3-(6-methylsulfonylpyridin-3-yl)phenyl]-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 51** in methylene chloride (30ml/mmol) was added urea-hydrogen peroxide (8eq) and the resulting mixture was cooled to 0°C. Trifluoroacetic acid (4.7eq) was added and the mixture was warmed to room temperature as a solution was obtained. After 18 hours, more urea-hydrogen peroxide (2.6eq) and trifluoroacetic acid (2eq) were added and stirring was continued for 2 hours. The mixture was quenched with saturated aqueous sodium metabisulfite, diluted with methylene chloride and the organic phase was washed with 1N aqueous HCl, then brine and water, dried and evaporated. The crude product was chromatographed on silica gel eluting with 40% toluene in acetone to afford the title compound as a solid.




<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.66 (m, 2H), 0.85 (m, 2H), 2.97 (m, 1H), 3.52 (s, 3H), 7.48 (m, 1H), 7.58-7.65 (m, 3H), 7.72-7.78 (m, 2H), 8.15 (d, 1H), 8.54 (s, 1H), 8.68 (brs, 1H), 8.81 (d, 1H), 9.01 (s, 1H), 9.71 (br, NH).

5

**EXAMPLE 76**

**N-Cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide**



10

Following the procedure of **Step 2** of **Example 30**, but substituting N-cyclopropyl-1-{5-bromo-3-[6-(1-hydroxy-1-methylethyl)pyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide from **Example 47** for 5-bromo-2-(1-hydroxy-1-methylethyl)pyridine, the title compound was obtained as a solid.


<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.71 (m, 2H), 0.90 (m, 2H), 1.75 (s, 6H), 3.02 (m, 1H), 7.48-7.60 (m, 5H), 7.73 (s, 1H), 7.88 (s, 1H), 8.52 (s, 1H), 8.72 (m, 1H), 8.84 (dd, 1H), 9.04 (s, 1H), 9.71 (br, NH).

15

20

**EXAMPLE 77**

**N-Cyclopropyl-1-{3-[6-(1,2-dihydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.**



**Step 1: 5-Bromo-2-(1-methylvinyl)pyridine N-oxide**

A mixture of 5-bromo-2-(1-hydroxy-1-methylethyl) pyridine N-oxide from step 2 of example 30 (1.29g) and 25% aqueous sulfuric acid was heated at 130°C for 2 days. After cooling, the mixture was made slightly basic using 10N aqueous sodium hydroxide and partitioned between ethyl acetate and water. The crude product from evaporation of the organic phase was used as such in step 2.

**Step 2: 5-bromo-2-(1,2-dihydroxy-1-methylethyl) pyridine N-oxide**

The crude product from step 1 was dissolved in a 3 : 1 mixture of acetone and water (16mL) and 4-methylmorpholine N-oxide (1g) and potassium osmate dihydrate (90mg) were added. The resulting mixture was stirred at room temperature for 3 days then excess solid sodium bisulfite was added and the mixture was evaporated. The residue was diluted with methylene chloride and filtered. The filtrate was evaporated and the residue chromatographed on silica gel eluting with ethyl acetate to afford the title compound as a white solid.

**Step 3: N-Cyclopropyl-1-{3-[6-(1,2-dihydroxy-1-methylethyl)-1-oxidopyridin-3-yl]phenyl}-1,4-dihydro[1,8]naphthyridin-4-one-3-carboxamide.**

Following the procedure of step 2 of example 32 but substituting 5-bromo-2-(1,2-dihydroxy-1-methylethyl) pyridine N-oxide from step 2 for 3-bromo-5-methylsulfonylpyridine the title compound was obtained as a white solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 0.66 (m, 2H), 0.85 (m, 2H), 1.61 (s, 3H), 2.78 (m, 1H, OH), 2.97 (m, 1H), 3.90 (m, 1H), 3.97 (m, 1H), 7.48 (m, 1H), 7.53 (m, 2H), 7.60 (m, 2H), 7.69-7.72 (m, 2H), 7.92 (s, 1H, OH), 8.49 (s, 1H), 8.68 (m, 1H), 8.80 (dd, 1H), 9.02 (s, 1H), 9.73 (br, NH).


5

Other variations or modifications, which will be obvious to those skilled in the art, are within the scope and teachings of this invention. This invention is not to be limited except as set forth in the following claims.

10

## WHAT IS CLAIMED IS:

1. A compound represented by Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein

Ar is phenyl, pyridyl, pyrimidyl, indolyl, quinolinyl, thienyl, pyridonyl, oxazolyl, oxadiazolyl, thiadiazolyl, or imidazolyl; or oxides thereof when Ar is a heteroaryl;

R is H or -C<sub>1-6</sub>alkyl;

R<sup>1</sup> is H, or -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkoxy, -C<sub>2-6</sub>alkenyl, -C<sub>3-6</sub>alkynyl, heteroaryl, or heterocycle group, optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents;

R<sup>2</sup> is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>3-6</sub>cycloalkyl, -C<sub>1-6</sub>alkyl(C<sub>3-6</sub>cycloalkyl)(C<sub>3-6</sub>cycloalkyl), -C<sub>1-6</sub>alkoxy, phenyl, heteroaryl, heterocycle, amino, -C(O)-C<sub>1-6</sub>alkyl, -C(O)-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl(=N-OH), -C(N=NOH)C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl(oxy)C<sub>1-6</sub>alkyl-phenyl, -SO<sub>n</sub>NH(C<sub>0-6</sub>alkyl), or -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), wherein the phenyl, heteroaryl or heterocycle is optionally substituted with halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, hydroxy, amino, or -C(O)-O-C<sub>1-6</sub>alkyl, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents;

n is 0, 1, or 2;

R<sup>3</sup> is H, OH, amine, halogen or C<sub>1-6</sub>alkyl, wherein the alkyl is optionally substituted with 1-6 independent halogen, OH, or amine substituents; and

R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is H, halogen, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, or amine, and any alkyl is optionally substituted with 1-6 independent halogen or -OH substituents.

5                    2.        The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

Ar is phenyl.

10                   3.        The compound according to claim 2, or a pharmaceutically acceptable salt thereof, wherein

R<sup>1</sup> is -C<sub>1-6</sub>alkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

15                   4.        The compound according to claim 2, or a pharmaceutically acceptable salt thereof, wherein

R<sup>1</sup> is -C<sub>3-6</sub>cycloalkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

20                   5.        The compound according to claim 2, or a pharmaceutically acceptable salt thereof, wherein  
R<sup>1</sup> is a pyridyl group optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

30                   6. The compound according to claim 1, or a pharmaceutically acceptable salt, wherein

Ar is pyridyl, pyrimidyl, or oxide thereof.

7. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein

R is H; and

R<sup>1</sup> is H.

8. The compound according to claim 6, or a pharmaceutically acceptable salt, wherein

5 R<sup>1</sup> is -C<sub>1-6</sub>alkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

10 9. The compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein

R<sup>1</sup> is -C<sub>3-6</sub>cycloalkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

15 10. The compound according to claim 6, or a pharmaceutically acceptable salt, wherein

R<sup>1</sup> is pyridyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

20 11. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

Ar is indolyl, quinoliny, or oxide thereof.

25 12. The compound according to claim 11, or a pharmaceutically acceptable salt thereof, wherein

R<sup>1</sup> is -C<sub>1-6</sub>alkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.

30 13. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

Ar is thienyl.

14. The compound according to claim 13, or a pharmaceutically acceptable salt thereof, wherein

5  $R^1$  is -C<sub>3-6</sub>cycloalkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.


15. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein




10 Ar is pyridonyl.

16. The compound according to claim 15, or a pharmaceutically acceptable salt thereof, wherein




15  $R^1$  is -C<sub>3-6</sub>cycloalkyl optionally substituted with 1-3 independent -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkoxy, OH, amino, -(C<sub>0-6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1-6</sub>alkyl), nitro, CN, =N-O-C<sub>1-6</sub>alkyl, -O-N=C<sub>1-6</sub>alkyl, or halogen substituents.




17. The compound according to claim 1, comprising




































or a pharmaceutically acceptable salt thereof.




5




18. The compound according to claim 1, comprising



















































































or a pharmaceutically acceptable salt thereof.


19. The compound according to claim 1, comprising



5








or a pharmaceutically acceptable salt thereof.

20. The compound according to claim 1, comprising



or a pharmaceutically acceptable salt thereof.

21. A pharmaceutical composition comprising  
a therapeutically effective amount of a compound of Formula (I) as  
defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt thereof; and  
a pharmaceutically acceptable carrier.

22. The pharmaceutical composition according to claim 21, further  
comprising a Leukotriene receptor antagonist, a Leukotriene biosynthesis inhibitor, an  
M2/M3 antagonist, a corticosteroid, an H1 receptor antagonist or a beta 2  
adrenoceptor agonist.

23. The pharmaceutical composition according to claim 21, further comprising a COX-2 selective inhibitor, a statin, or an NSAID.

5                   24. A method of treatment or prevention of asthma, chronic  
bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma,  
psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock  
(and associated conditions such as laminitis and colic in horses), septic shock,  
ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain,  
10 inflammatory arthritis, osteoporosis, chronic glomerulonephritis, atopic dermatitis,  
urticaria, adult respiratory distress syndrome, infant respiratory distress syndrome,  
chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis,  
allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, atherosclerosis,  
neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis,  
15 transplant rejection and graft versus host disease, hypersecretion of gastric acid,  
bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-  
mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting,  
depression, memory impairment, monopolar depression, acute and chronic  
neurodegenerative disorders with inflammatory components, Parkinson disease,  
20 Alzheimer's disease, spinal cord trauma, head injury, multiple sclerosis, tumour  
growth and cancerous invasion of normal tissues comprising the step of administering  
a therapeutically effective amount, or a prophylactically effective amount, of the  
compound according to claim 1 or a pharmaceutically acceptable salt thereof.

25. Use of a compound of Formula (I), as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment or prevention of the diseases or ailments set forth in Claim 24.

5

26. A compound of Formula (I), as defined in any one of Claims 1 to 20, or a pharmaceutically acceptable salt thereof, for use in therapeutic inhibitors of phosphodiesterase -4.

10

27. A phosphodiesterase-4 inhibitor pharmaceutical composition comprising an acceptable, phosphodiesterase-4 inhibiting amount of a compound of Formula (I), as defined in any one of Claims 1 to 20, in association with a pharmaceutically acceptable carrier.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/02/00746

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/04 A61K31/435 A61K31/495 A61P11/06 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, PAJ, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 07704 A (SUNTORY LTD ;INOUE HIDEKAZU (JP); HAYASHI YASUHIRO (JP); SHIMAMOTO) 18 February 1999 (1999-02-18) cited in the application page-7, line 21 - page 58 line 2 page 2, line 31 -page 4, line 29; claim 1; examples 144,146,148,149	1-27
Y	EP 0 978 516 A (SUNTORY LTD) 9 February 2000 (2000-02-09) page 3, line 16 -page 3, line 41; claim 1	1-27
A	WO 97 04775 A (CHIROSCIENCE LTD) 13 February 1997 (1997-02-13) page 4, line 1 -page 5, line 11; claim 1 -/--	1-27

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

\* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\* document member of the same patent family

Date of the actual completion of the international search

24 September 2002

Date of mailing of the international search report

04/10/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Wörth, C

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 02/00746

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 492 702 A (SHERLOCK MARGARET H) 8 January 1985 (1985-01-08) claim 28	1-27

## INTERNATIONAL SEARCH REPORT

on patent family members

Inter. Application No.

PCT/JP 02/00746

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9907704	A	18-02-1999	AU 8560798 A	01-03-1999
			CN 1239478 T	22-12-1999
			EP 0958297 A1	24-11-1999
			HU 0001054 A2	28-10-2000
			WO 9907704 A1	18-02-1999
			JP 11106385 A	20-04-1999
			US 6297248 B1	02-10-2001
			US 2002006935 A1	17-01-2002
EP 0978516	A	09-02-2000	AU 2185699 A	16-08-1999
			EP 0978516 A1	09-02-2000
			HU 0002353 A2	28-05-2001
			US 6331548 B1	18-12-2001
			CN 1256693 T	14-06-2000
			WO 9938867 A1	05-08-1999
WO 9704775	A	13-02-1997	AU 695132 B2	06-08-1998
			AU 6626696 A	26-02-1997
			CA 2225555 A1	13-02-1997
			EP 0841927 A1	20-05-1998
			WO 9704775 A1	13-02-1997
			JP 11510156 T	07-09-1999
			US 5753666 A	19-05-1998
			ZA 9606600 A	04-08-1997
			AU 696390 B2	10-09-1998
			AU 6626396 A	26-02-1997
			CA 2225552 A1	13-02-1997
			EP 0841929 A1	20-05-1998
			WO 9704779 A1	13-02-1997
			JP 11513021 T	09-11-1999
			US 5891878 A	06-04-1999
US 4492702	A	08-01-1985	US 4452800 A	05-06-1984
			AT 24727 T	15-01-1987
			AU 553958 B2	31-07-1986
			AU 1392783 A	03-11-1983
			CA 1198109 A1	17-12-1985
			DE 3368929 D1	12-02-1987
			DK 181983 A	27-10-1983
			EP 0092786 A2	02-11-1983
			HK 20590 A	23-03-1990
			IE 55113 B1	06-06-1990
			JP 5068476 B	29-09-1993
			JP 58194887 A	12-11-1983
			NZ 204004 A	14-03-1986
			PH 21641 A	13-01-1988
			PT 76596 A ,B	01-05-1983
			SG 76889 G	20-04-1990
			ZA 8302903 A	25-01-1984
			FI 833137 A ,B,	04-05-1984
			GR 79325 A1	22-10-1984
			HU 189715 B	28-07-1986
			KR 8904662 B1	24-11-1989
			NO 832534 A ,B,	04-05-1984
			OA 7522 A	31-03-1985
			US 4551463 A	05-11-1985
			US 4628055 A	09-12-1986